

Automorphic period and the central value of Rankin-Selberg L-function

Wei Zhang

August 31, 2012

Abstract

We prove a refinement of the global Gan–Gross–Prasad conjecture essentially proposed by Ichino-Ikeda and N. Harris for unitary groups under some local conditions. We need to assume some expected properties of L-packets and some part of the local Gan-Gross-Prasad conjecture.

Contents

1	Introduction	1
2	Measures	10
3	Explicit local factorization of some periods	13
4	Relative trace formulae of Jacquet and Rallis	21
5	The totally definite case	33
6	Harmonic analysis on Lie algebra	34
7	Smoothing local periods	43
8	Local character expansion in the general linear case	52
9	Local character expansion in the unitary case	69

1 Introduction

In this article, as a sequel of [50], we prove a conjectural refinement of the global Gan–Gross–Prasad conjecture ([6]) for unitary groups under some local conditions. In a influential

paper [23], Ichino and Ikeda first formulated the refinement for orthogonal groups. After the Ichino–Ikeda formulation, N. Harris considered the case of unitary groups in his Ph.D Thesis at UCSD ([21]). Our version of the refinement is slightly different from theirs at the archimedean place.

The conjecture of Ichino–Ikeda and N. Harris. We now recall the conjectural refinement. Let E/F be a quadratic extension of number fields with adeles denoted by $\mathbb{A} = \mathbb{A}_F$ and $\mathbb{A}_E$ respectively. Let V be a Hermitian space of dimension $n + 1$ and W a (non-degenerate) subspace of codimension one. Denote the unitary groups by $U(V)$ and $U(W)$ respectively. Let $G = U(W) \times U(V)$ be the product and H the diagonal embedding of $U(W)$ into G . Denote $G' = \mathrm{GL}_{n,E} \times \mathrm{GL}_{n+1,E}$ as an algebraic group over E . Let $\pi = \pi_n \otimes \pi_{n+1}$ be a cuspidal automorphic representation of $G(\mathbb{A})$ and let $\pi_{i,E}$ be the base change of π_i to $\mathrm{GL}_i(\mathbb{A}_E)$, $i = n, n + 1$. Denote by $L(s, \pi_E)$ the Rankin–Selberg convolution L-function $L(s, \pi_{n,E} \times \pi_{n+1,E})$. It is known to be the same as the one defined by the Langlands–Shahidi method. The reader may consult the introduction of [10] for an overview of the study of the central value $L(1/2, \pi_{n,E} \times \pi_{n+1,E})$. We also consider the adjoint L-function of π

$$L(s, \pi, \mathrm{Ad}) = L(s, \pi_n, \mathrm{Ad})L(s, \pi_{n+1}, \mathrm{Ad}).$$

Denote the constant

$$\Delta_{n+1} = \prod_{i=1}^{n+1} L(i, \eta^i) = L(1, \eta)L(2, 1_F)L(3, \eta) \cdots L(n + 1, \eta^{n+1}),$$

where η is the quadratic character of $\mathbb{A}^\times$ associated to E/F by class field theory. Note that here $\Delta_{n+1} = L(M^\vee(1))$ where $M^\vee$ is the motive dual to the motive M associated to the reductive group $U(n)$ defined by Gross ([13]). We will be interested in the following combination of L-functions

$$(1.1) \quad \mathcal{L}(s, \pi) = \Delta_{n+1} \frac{L(s, \pi_E)}{L(s + \frac{1}{2}, \pi, \mathrm{Ad})}.$$

We also write $\mathcal{L}(s, \pi_v)$ the local component at v .

Let $[H]$ denote the quotient $H(F) \backslash H(\mathbb{A})$ and similarly for G . We endow $H(\mathbb{A})$ ($G(\mathbb{A})$, resp.) with their Tamagawa measures ¹ and $[H]$ ($[G]$, resp.) with the quotient measure by the counting measure on $H(F)$ ($G(F)$, resp.), cf. §2. In [6], Gan, Gross and Prasad propose to study an automorphic period integral

$$\mathcal{P}(\phi) = \mathcal{P}_H(\phi) := \int_{[H]} \phi(h) dh, \quad \phi \in \pi.$$

¹Since the unitary group H has nontrivial central torus, we need to introduce a convergence factor: $dh = L(1, \eta)^{-1} \prod_v L(1, \eta_v) |\omega|_v$ for a nonzero invariant differential ω on H . Similarly for G .

They conjecture that the non-vanishing of the linear functional $\mathcal{P}$ on π (possibly by switching to another member in the Vogan L-packet of π) is equivalent to the non-vanishing of the central value $L(\frac{1}{2}, \pi_E)$ of the Rankin-Selberg L-function. This conjectural equivalence is proved for π satisfying some local conditions in our previous paper ([50]). One direction of the equivalence was also proved by Ginzburg–Jiang–Rallis (cf. [10], [11]).

For arithmetic application, it is necessary to have more precise relation between the automorphic period integral $\mathcal{P}$ and the central L-value. To state the precise refinement of the Gan-Gross-Prasad conjecture, we need to introduce more notation. Let $\langle \cdot, \cdot \rangle_{Pet}$ be the Peterson inner product

$$\langle \phi, \varphi \rangle_{Pet} = \int_{[G]} \phi(g) \overline{\varphi}(g) dg, \quad \phi, \varphi \in \pi.$$

Fix an decomposition as a product

$$\langle \cdot, \cdot \rangle_{Pet} = \prod_v \langle \cdot, \cdot \rangle_v$$

under the decomposition $\pi = \otimes \pi_v$. In this way we fix an invariant inner product on π_v . Ichino and Ikeda first consider the following integration of matrix coefficient: for $\phi_v, \varphi_v \in \pi_v$, we define

$$\alpha_v(\phi_v, \varphi_v) = \int_{H_v} \langle \pi_v(h) \phi_v, \varphi_v \rangle_v dh, \quad H_v = H(F_v).$$

It has the following nice properties:

1. If π_v is tempered, it converges absolutely and $\alpha(\phi_v, \phi_v) \geq 0$.
2. When π_v is unramified², and the vectors ϕ, φ are fixed by K_v such that $\langle \phi_v, \varphi_v \rangle_v = 1$, we have

$$\alpha_v(\phi_v, \varphi_v) = \mathcal{L}(\frac{1}{2}, \pi_v) \cdot \text{vol}(H(\mathcal{O}_v)).$$

3. If $\text{Hom}_{H(F_v)}(\pi_v, \mathbb{C}) \neq 0$, then the form α_v does not vanish identically.

The first two are proved by Ichino and Ikeda (N. Harris in the unitary case). The third property is conjectured by them and proved by Sakellaridis and Venkatesh ([34]) in more general setting. Waldspurger also proved the third property in the orthogonal case. Because of the second property, we normalize the form α_v

$$\alpha_v^{\natural}(\phi_v, \varphi_v) = \frac{1}{\mathcal{L}(\frac{1}{2}, \pi_v)} \int_{H_v} \langle \pi_v(h) \phi_v, \varphi_v \rangle_v dh.$$

²For a place v we say that π_v is unramified if the quadratic extension E/F is unramified at v , the group $G(F_v)$ has a hyperspecial subgroup $K_v = G(\mathcal{O}_v)$ and π_v has a nonzero K_v -fixed vector.

We are now ready to state the conjecture of Ichino-Ikeda and N. Harris (cf. [23], [21]) that refines the global Gan–Gross–Prasad conjecture for unitary groups. Assume that the measure on $H(\mathbb{A})$ and the measures on $H(F_v)$ defining α_v satisfy:

$$dh = \prod_v dh_v.$$

Conjecture 1.1. *Assume that π is tempered. For any decomposable vector $\phi = \otimes \phi_v \in \pi = \otimes \pi_v$, we have*

$$(1.2) \quad \frac{|\mathcal{P}(\phi)|^2}{\langle \phi, \phi \rangle_{Pet}} = \frac{1}{|S_\pi|} \mathcal{L}\left(\frac{1}{2}, \pi\right) \prod_v \frac{\alpha_v^{\natural}(\phi_v, \phi_v)}{\langle \phi_v, \phi_v \rangle_v},$$

where S_π is a finite elementary 2-group: the component group associated to the L-parameter of $\pi = \pi_n \otimes \pi_{n+1}$.

The conjectural formula of this kind goes back to the celebrated work of Waldspurger ([38]) for the central values of L-functions of $\mathrm{GL}(2)$ (more or less equivalent to the case $U(1) \times U(2)$ in the unitary setting). An arithmetic geometric version generalizing the formula of Gross–Zagier and S. Zhang ([16], [43]) is also formulated in [48]. More explicit formulae were obtained by Gross ([12]), S. Zhang ([47]) and many others. The formula of Waldspurger and the formula of Gross–Zagier and S. Zhang ([16], [45], [46], [43]) play an important role in the spectacular development in application to the Birch and Swinnerton-Dyer conjecture for elliptic curves in the past thirty years. More recently, Y. Tian ([37]) applies both formulae together to a classical Diophantine question and proves the infinitudes of square-free congruent numbers with an arbitrary number of primes factors.

The refined global conjecture for $\mathrm{SO}(3) \times \mathrm{SO}(4)$, concerning “the triple product L-function”, was established after the work by Garret [8], Piatetski-Shapiro–Rallis, Harris–Kudla [19], Gross–Kudla, and Ichino [22]. Recently Gan and Ichino ([7]) established some new cases for $\mathrm{SO}(4) \times \mathrm{SO}(5)$ (for endoscopic L-packets on $\mathrm{SO}(5)$). All of the known cases utilize the theta correspondence in an ingenious way.

The formula of Waldspurger was also reproved by Jacquet and Jacquet-Chen ([3]) using relative trace formulae. Our method is more close to theirs.

Main results. We now state our main result. Throughout this paper, we assume *the expected properties the L-packets of unitary groups (for all Hermitian spaces W, V), analogous to the work of Arthur on orthogonal groups* (cf. [31], [41] for the progress towards this end). Moreover we need to assume a part of the local Gan-Gross-Prasad conjecture in the unitary case for *all local fields of characteristic zero: in a tempered (Vogan) L-packet there is at most one member that admits a nonzero invariant linear form*. A proof of this part (in fact an even stronger result) for p -adic fields is recently posted by Beuzart-Plessis ([2, Theorem 1]); the local conjecture in the orthogonal case for p -adic fields has earlier been proved by Waldspurger.

We also assume that the fundamental lemma of Jacquet–Rallis holds for all p -adic field. When $n = 2$ and $p > 2$, this was verified by the author by brutal computation. In [44] and its appendix by J. Gordon, the fundamental lemma is proved when p is larger than a constant $c(n)$. It is plausible that one may deduce the general case from their results. Without assuming this, in the theorem below we will need to impose one more condition on the quadratic extension E/F : it is split for all non-archimedean place with residue characteristic $\leq c(n)$.

At a split place v , we may also define an alternative normalization via the Whittaker model:

$$\alpha'_v(\phi_v, \varphi_v) = \lambda_v(W_{\phi_v}) \overline{\lambda_v(W_{\varphi_v})}.$$

See §3 for the precise definition of λ_v in terms of the Whittaker model and normalization of the inner product on the Whittaker model. When v is non-archimedean, we will show that

$$\alpha_v = \alpha'_v.$$

It is expected to be true for archimedean places, too. Also we set $\alpha'_v = \alpha_v$ for non-split v . We again normalize

$$\alpha_v^{\natural}(\phi_v, \varphi_v) = \frac{1}{\mathcal{L}(\frac{1}{2}, \pi_v)} \alpha'_v(\phi_v, \varphi_v).$$

Theorem 1.2. *Let π be tempered cuspidal automorphic representation of $G(\mathbb{A})$. Denote Σ by the set of non-split places v of F where π_v is not unramified. Assume that*

- (i) *The local components π_v are supercuspidal for at least one split place.*
- (ii) *If $v \in \Sigma$, then either H_v is compact or π_v is supercuspidal.*

Then we have results for the following two (extreme) cases according to $G(\mathbb{R})$:

- (1) *The totally split case: if every archimedean place v of F is split in the extension E/F (i.e., $G_{F_\infty} \simeq (\mathrm{GL}_n \times \mathrm{GL}_{n+1})_{F_\infty}$), then we have*

$$\frac{|\mathcal{P}(\phi)|^2}{\langle \phi, \phi \rangle_{Pet}} = 2^{-2} \mathcal{L}\left(\frac{1}{2}, \pi\right) \prod_v \frac{\alpha_v^{\natural}(\phi_v, \phi_v)}{\langle \phi_v, \phi_v \rangle_v}.$$

- (2) *The totally definite case: if $G(F_\infty)$ is compact where $F_\infty = \prod_{v|\infty} F_v$ (in particular, E is a CM extension of a totally real field F), then the same equality holds up to a non-zero constant c_{π_∞} depending only on the archimedean component π_∞ of π :*

$$\frac{|\mathcal{P}(\phi)|^2}{\langle \phi, \phi \rangle_{Pet}} = c_{\pi_\infty} 2^{-2} \mathcal{L}\left(\frac{1}{2}, \pi\right) \prod_v \frac{\alpha_v^{\natural}(\phi_v, \phi_v)}{\langle \phi_v, \phi_v \rangle_v}.$$

Under our assumptions (i), the automorphic representation π has cuspidal base change to the general linear group and hence

$$|S_\pi| = |S_{\pi_n}| \cdot |S_{\pi_{n+1}}| = 4.$$

The condition (i) is due to the fact that currently we do not have a complete spectral decomposition of the relative trace formulae of Jacquet and Rallis. The condition (ii) seems to be only a technical restriction for our approach and will be discussed in §9. The restriction for the archimedean place is due to the lack of the existence of smooth transfer at archimedean places (cf. §5). However, even if we assume the existence of smooth transfer, it is probably substantially more technical to remove the restriction in the archimedean case (say, to evaluate the constant c_{π_∞}).

We also make a local conjecture (Conjecture 4.3) for each place v . Assuming this conjecture, we may improve Theorem 1.2 to only assume that the base change π_E is cuspidal.

Remark 1. At a split place, in practical computation it seems easier to use the normalization via Whittaker model instead of the matrix coefficient integration, for example, the archimedean computation by Watson for the triple product L-function ([42]).

Remark 2. In our previous paper [50], we proved the equivalence of non-vanishing of the two sides of the equality (in more generality). It is worth mentioning that the current result relies on this equivalence; indeed the equivalence shows that if LHS is zero, so is the RHS. What the current paper actually deals with is the case when the LHS is non-zero.

Remark 3. Note that for a non-archimedean place v , H_v is possibly compact only when $n \leq 2$. When $n = 1$, H_v is always compact for a non-split v . This compact case already occurred in [3].

Some applications. We have the following application to the positivity of some L-values. The positivity is also predicted by the grand Riemann hypothesis. Lapid has obtained a more general result for Rankin–Selberg central L-values via a different method ([30], cf. also [29] for the positivity of central values of L-functions of symplectic type).

Theorem 1.3. *Assume that π is as above and E/F is split at all archimedean places. Then we have*

$$L(\pi_E, \frac{1}{2}) \geq 0.$$

Proof. It suffices to show this when $L(\pi_E, \frac{1}{2}) \neq 0$. Then by [50], there exists π' in the same L-packet of π such that the period $\mathcal{P}$ on π' does not vanish. By replacing π by π' , we may assume that the space $\text{Hom}_{U(W)(F_v)}(\pi_v, \mathbb{C})$ does not vanish for every v . Then the local terms α'_v do not vanish. Now the positivity follows from the fact that the α'_v are all positive definite, and the other L-values appearing in $\mathcal{L}(\pi_E, 1/2)$ except $L(\pi_E, \frac{1}{2})$ are all positive. $\square$

Remark 4. As another application, M. Harris showed that the Ichino-Ikeda conjecture would imply the algebraicity of the L-value $\mathcal{L}(1/2, \pi)$ up to some simple constant when $G(F_\infty)$ is compact and $\alpha_\infty \neq 0$ (cf. [20, §4.1]).

Outline of proof. We now sketch the main ideas of the proof. First of all, by the multiplicity one result ([1],[36]), we know a priori that there is a constant denoted by $\mathcal{C}_\pi$ depending on π such that for all decomposable $\phi, \varphi \in \pi$:

$$(1.3) \quad \mathcal{P}(\phi) \overline{\mathcal{P}(\varphi)} = \mathcal{C}_\pi \prod_v \alpha_v^\natural(\phi_v, \varphi_v).$$

Instead of working with individual ϕ as in the conjecture, we switch our angle of view to working with a distribution attached to π :

$$(1.4) \quad J_\pi(f) := \sum_{\phi} \mathcal{P}(\pi(f)\phi) \overline{\mathcal{P}(\phi)}, \quad f \in \mathcal{C}_c^\infty(G(\mathbb{A})),$$

where the sum of ϕ is over an orthonormal basis of π . In this article, we call this distribution a (*global*) *spherical character* following the names of many of its early analogues (cf. [32] etc.). We may accordingly define a *local spherical character*:

$$(1.5) \quad J_{\pi_v}^\natural(f_v) := \sum_{\phi_v} \alpha_v^\natural(\pi_v(f_v)\phi_v, \phi_v), \quad f_v \in \mathcal{C}_c^\infty(G(F_v)),$$

where the sum of ϕ_v is over an orthonormal basis of π_v . By (1.3), we clearly have for decomposable $f = \bigotimes_v f_v$:

$$(1.6) \quad J_\pi(f) = \mathcal{C}_\pi \prod_v J_{\pi_v}^\natural(f_v),$$

where in the product of RHS, for a given π and f , the local term $J_{\pi_v}^\natural(f_v) = 1$ for all but finitely many v . Then we have the following consequence of Conjecture 1.1.

Conjecture 1.4. *For all $f = \bigotimes_v f_v \in \mathcal{C}_c^\infty(G(\mathbb{A}))$, we have*

$$J_\pi(f) = \mathcal{L}(\pi, 1/2) \prod_v J_{\pi_v}^\natural(f_v).$$

In fact, Conjecture 1.4 also implies, hence is equivalent to Conjecture 1.1. To see this, we note that the following are equivalent: (1) $\text{Hom}(\pi_v, \mathbb{C}) \neq 0$, (2) $\alpha_v \neq 0$, (3) *the distribution $J_{\pi_v}^\natural$ does not vanish*.³ Hence, Conjecture 1.1 holds if for some v the linear form α_v vanishes. Now assume that for all v , the linear forms α_v do not vanish. Then the distributions J_{π_v} do not vanish. Then by Conjecture 1.4, the constant $\mathcal{C}_\pi$ must be $\mathcal{L}(\pi, 1/2)$ which implies Conjecture 1.1. Similarly, Theorem 1.2 follows from similar conjecture as 1.4 where one replaces α_v by α'_v when defining J_{π_v} .

Moreover, there is a parallel question on general linear groups and it is essentially reduced to the celebrated theory of “Rankin–Selberg convolution” ([25]). Therefore, a natural idea

³It is clear that (2) is equivalent to (3). The equivalence of (1) and (2) follows from the third property of α_v listed earlier.

is to transfer the question from unitary groups to general linear groups via (quadratic) base change from unitary groups to general linear groups. We are therefore lead to comparing the two period integrals via the relative trace formulae of Jacquet and Rallis ([26]).

In the general linear case, there is a decomposition of a global spherical character into local ones, analogous to Conjecture 1.4. But this time one may prove it without difficulty (indeed almost tautological). Hence, to deduce the Conjecture 1.4, it suffices to compare the two local spherical characters. Moreover, since we only need to find the constant $\mathcal{C}_\pi$, we may just choose some special test functions f (as long as the RHS does not vanish for our choice). Therefore the main innovation of this paper is a formula for the local spherical character evaluated at some special test functions. The formula can be viewed as a truncated *local expansion of the local spherical character* that is analogous to the local expansion of characters due to Harish-Chandra. The result may be of independent interest in view of local harmonic analysis in the relative setting.

For comparison, let us recall briefly the result of Harish-Chandra. Let F be a p -adic fields, G the F -points of a connected reductive group, and $\mathfrak{g}$ the Lie algebra of G . Let $\mathcal{N}$ be the nilpotent cone of $\mathfrak{g}$ and denote by $\mathcal{N}/G$ the set of G -conjugacy classes in $\mathcal{N}$, which is a finite set. Let $\mu_{\mathcal{O}}$ be the nilpotent orbital integral associated to $\mathcal{O} \in \mathcal{N}/G$. The exponential map defines an homeomorphism $\exp : \omega \rightarrow \Omega$ where ω (Ω , resp.) is some neighborhood of 0 in $\mathfrak{g}$ (G , resp.). Let π be an irreducible admissible representation of G . Then Harish-Chandra showed that there are constants $c_{\mathcal{O}}(\pi)$ indexed by $\mathcal{O} \in \mathcal{N}/G$ such that when Ω is sufficiently small, for all f supported in Ω :

$$(1.7) \quad \mathrm{tr}(\pi(f)) = \sum_{\mathcal{O} \in \mathcal{N}/G} c_{\mathcal{O}}(\pi) \mu_{\mathcal{O}}(\hat{f}_{\mathfrak{h}}).$$

Here $f_{\mathfrak{h}}$ is the function on ω via the homeomorphism, and $\hat{f}_{\mathfrak{h}}$ its Fourier transform. The constants $c_{\mathcal{O}}(\pi)$ contains important information of π . For example, there is a distinguished nilpotent conjugacy class, namely the conjugacy of $0 \in \mathcal{N}$. If π is a discrete series representation, the constant $c_{\{0\}}(\pi)$ is essentially the formal degree of π provided an appropriate choice of Haar measure.

Now we return to our local setting for a non-split non-archimedean place v of F . We suppress the subscript v . We consider the local spherical character in the general linear case. Let $G' := \mathrm{Res}_{E/F}(GL_n \times GL_{n+1})$ and let Π be irreducible unitary generic representations of $G'(F)$. Then the local spherical character I_{Π} (cf. (3.17)) defines a distribution on $G'(F)$ with certain invariance property. These distributions are related to distributions on the F -vector space:

$$\mathfrak{s}_{n+1} = \{X \in M_{n+1}(E) | X + \overline{X} = 0\}.$$

The group $GL_{n+1}(F)$ acts on $\mathfrak{s}_{n+1}$ by conjugation and we will be interested in the restriction of the action to the subgroup $GL_n(F)$ (as a factor of the Levi of the parabolic of $(n, 1)$ -type). We let ω be a small neighborhood of 0 in the F -vector space $\mathfrak{s}_{n+1}$. Then we have a natural way to pull back a function f' on a small neighborhood of 1 in G' to a function denoted by $f'_{\mathfrak{h}}$ on ω (cf. §8 for the precise definition). It is tempting to guess that there exists an analogous

expansion of I_Π in terms of the (relative to $GL_n(F)$ -action) unipotent orbital integrals on $\mathfrak{s}_{n+1}$.⁴ However, so far there are some difficulty. For example, when $n \geq 2$ there are infinitely many $GL_n(F)$ -nilpotent orbits in $\mathfrak{s}_{n+1}$ and these nilpotent orbital integrals often need to be regularized. We then define a subspace of *admissible functions* on ω . The precise definition is very technical. We expect that admissible functions have vanishing nilpotent orbital integrals (however, generally not even defined so far), except for one of the two regular unipotent orbits denoted by ξ_- . An expansion such as (1.7) of $I_\Pi(f)$ would then tell us that there should be only one term left, corresponding to the regular unipotent orbit ξ_- . Though it seems challenging to prove something such as (1.7) in our setting, we nevertheless manage to establish a truncated version (see Theorem 8.4 for the detail):

Theorem 1.5. *Let Π be irreducible unitary generic representations of $G'(F)$. Then there is a sufficiently small ω such that for every admissible function f' ,*

$$I_\Pi(f') = (*)\mu_{\xi_-}(\widehat{f'_\mathfrak{h}}) \neq 0,$$

where $(*)$ is an explicit non-zero constant independent of Π (up to its central character).

We have a similar result for a local spherical character J_π on the unitary group when either π is a supercuspidal representation or the group $U(W)$ is compact. See §9 for more details (Theorem 9.7). Then our main result follows from the local comparison of the two spherical characters (cf. §4 Conjecture 4.3).

The proof for the unitary case seems to be harder and needs the full strength of our previous results in the companion paper [50]. Namely we have to make use of the following results (cf. §9)

1. The existence of smooth transfer.
2. Compatibility of smooth transfer with Fourier transform.
3. Local (relative) trace formula on “Lie algebra”.

Note that the proof in [50] of the these ingredients are in the reverse order listed here.

Structure of this paper. After fixing some notations in §2, we review several global periods involving general linear groups in §3 and deduce the decomposition analogous to Conjecture 1.4. Then in §4 we recall the RTF of Jacquet and Rallis and reduce the question to a comparison of local spherical characters. Then we give the proof of Theorem 1.2 assuming a local result. In §5 we deal with the totally definite case (i.e., $G(\mathbb{R})$ compact). In §6 we prepare some (relative) harmonic analysis on Lie algebras. In §7 and §8, we prove the local character expansion for the general linear group. In §9 we show the local character expansion for the unitary group under some conditions, and complete the proof via the comparison of both characters.

⁴Relative to $GL_n(F)$ -action, an $X \in \mathfrak{s}_{n+1}$ is “nilpotent” if the closure of its $GL_n(F)$ -orbit contains zero.

Finally, we warn the reader of the change of measures: Only in the introduction, we use the Tamagawa measures associated to a differential form ω on H normalized by:

$$dh = L(1, \eta)^{-1} \prod_v L(1, \eta_v) |\omega|_v.$$

To have a natural local decomposition, below we will immediately switch to

$$dh = \prod_v dh_v, \quad dh_v = \prod_v L(1, \eta_v) |\omega|_v.$$

Another change comes when we go to the local setting where we even consider the unnormalized local measure

$$dh_v = |\omega|_v.$$

Similar warning applies to other groups such as G and the general linear groups.

Acknowledgement. The influence of Hervé Jacquet on this paper should be obvious to the reader. The author is grateful to S. Zhang and B. Gross for their constant support, to M. Harris, A. Ichino, D. Jiang, E. Lapid, C-P. Mok, Y. Tian, L. Xiao, X. Yuan for helpful comments. The author thanks the Morningside Center of Mathematics at the Chinese Academy of Sciences for their hospitality. The work is partially supported by NSF Grant DMS #1204365.

2 Measures

Measures related to general linear groups. We first list the main notations and conventions throughout this paper. We denote $H_n = \mathrm{GL}_n$, its standard Borel B_n with A_n the diagonal torus, N_n the unipotent radical. We denote by $N_{n,-}$ the opposite unipotent subgroup of H_n , and an open subvariety $H'_n = N_n A_n N_{n,-}$ of H_n (essentially the open cell of Bruhat decomposition). Their Lie algebras are denoted by $\mathfrak{h}_n, \mathfrak{n}_n$ etc. We denote $M_{n,m}(F)$ the F -vector space of all $n \times m$ -matrices with coefficients in F ; and if $n = m$ we write it as $M_n(F)$. Then it comes with the natural embedding $H_n \subset M_{n,n}$. We denote $e_n = (0, 0, \dots, 0, 1) \in M_{1,n}(F)$, $e_n^* \in M_{n,1}(F)$ the transpose of e_n .

We usually understand H_{n-1} as a subgroup of H_n via the block-diagonal embedding

$$H_{n-1} \ni h \mapsto \begin{pmatrix} h & \\ & 1 \end{pmatrix} \in H_n.$$

We thus have embeddings $\dots \subset H_{n-2} \subset H_{n-1} \subset H_n$. Similarly N_{n-1} as a subgroup of N_n etc..

For a quadratic extension E/F , we assume that $F = F[\tau]$ for $\tau = \sqrt{\delta}$. We write $E^\pm$ the F -vector space where the nontrivial Galois automorphism in $\mathrm{Gal}(E/F)$ acts by ± 1 . Locally we will fix an additive character $\psi = \psi_F$ of F and then define a character ψ_E of E by $\psi_E(z) = \psi(\frac{1}{2} \mathrm{tr}_{E/F} z)$ for the trace map $\mathrm{tr}_{E/F} : E \rightarrow F$. We also say that ψ is unramified if the largest ideal of F over which ψ is trivial is $\mathcal{O}_F$. Similarly for ψ_E . We will consider the self-dual measure on F , E^- and E . Note that if we use the isomorphism $F \simeq E^-$ by

$x \mapsto \sqrt{\delta}x$, then the measure on E^- is $|\delta|_F^{1/2} dx$ for the self-dual measure dx on E^- . Here our absolute values on F and E are normalized such that

$$d(ax) = |a|_F dx, x \in F$$

and similarly for E . On $M_n(E)$ there is a bilinear pairing given by $\text{tr}(XY)$ valued in E . We then have a Fourier transform for $\phi \in \mathcal{C}_c^\infty(M_n(E))$

$$\phi(X) := \int_{M_n(E)} \phi(Y) \psi_E(\text{tr}(XY)) dY,$$

for the self-dual measure on $M_n(E)$. Note that this is also the same measure by identifying $M_n(E)$ with E^{n^2} and using the self-dual measure on E .

On $F^\times$ we denote the *normalized* Tamagawa measure associated to the differential form $x^{-1}dx$:

$$d^\times x = \zeta_F(1) \frac{dx}{|x|_F},$$

and the *unnormalized* one:

$$d^*x = \frac{dx}{|x|_F}.$$

Similarly for $E^\times$. On $H_n(F)$ we will take the Haar measure

$$dg = \zeta_F(1) \frac{\prod_{ij} dx_{ij}}{|\det(g)|_F^n}, \quad g = (x_{ij}),$$

and similarly for $H_n(E)$ (where we replace $\zeta_F(1)$ by $\zeta_E(1)$). Sometimes we also shorten $|\det(g)|$ by $|g|$ if no confusion arises.

For $x \in M_{n,m}(F)$, we define

$$\|x\| = \max\{|x_{ij}|_F\}.$$

We will assign the measure on $N_n(F)$ the additive self-dual measure via the identification $N_n(F)$ with the F -vector space $F^{n(n-1)/2}$. Similarly we normalize the Haar measure on $A_n(F)$ by identifying it with $(F^\times)^n$.

We denote the modular character by

$$\delta_n(a) = \det(\text{Ad}(a) : \mathfrak{n}_n) = \prod_{i=1}^n |a_i|_F^{n+1-2i},$$

where $a = \text{diag}[a_1, a_2, \dots, a_n] \in A_n(F)$ acts on $\mathfrak{n}$ by $\text{Ad}(a)X = aXa^{-1}$ with unitary group $U(V)$. Similarly, we have $\delta_{n,E}$ if we replace F by E .

Now if F is a number field, we endow the group $H_n(\mathbb{A})$ with the product measure

$$dg = \prod_v dg_v.$$

Similarly for $H_n(\mathbb{A}_E)$ if E is a quadratic extension.

Measures related to unitary groups. Our method involves comparison of orbital integrals between the unitary and general linear cases, and between their Lie algebras, respectively. We thus need to choose compatible measures on them. Let θ be a Hermitian matrix of size $n+1$. Then we may and will view the group $U(\theta)(F)$ as the subgroup of $\mathrm{GL}_{n+1}(E)$ consisting of $g \in \mathrm{GL}_{n+1}(E)$ such that

$$\overline{g}^t \cdot \theta g \theta^{-1} = 1.$$

We may and will view the Lie algebra $\mathfrak{u}(\theta)$ of $U(\theta)$ as the subspace of $M_{n+1}(E)$ consisting of $X \in M_{n+1}(E)$ such that

$$\overline{X}^t + \theta X \theta^{-1} = 0.$$

We denote by $\mathfrak{u}(\theta)^\dagger$ a companion space where the last equality is replaced by

$$\overline{X}^t = \theta X \theta^{-1}.$$

For any number $\tau \in E$ such that $\overline{\tau} = -\tau \neq 0$, we have an isomorphism from $\mathfrak{u}(\theta)$ to $\mathfrak{u}(\theta)^\dagger$ mapping X to $\tau^{-1}X$.

We will need to consider the symmetric space $S_{n+1}(F) \simeq H_{n+1}(F) \backslash H_{n+1}(E)$, and identify it with the subspace of $\mathrm{GL}_{n+1}(E)$ consisting of $g \in \mathrm{GL}_{n+1}(E)$ such that

$$\overline{g}g = 1.$$

We also have the tangent space $\mathfrak{s} = \mathfrak{s}_{n+1}$ of $1 \in S_{n+1}$, which we call the *Lie algebra of $S_{n+1}(F)$* . Viewed as a subspace of $M_{n+1}(E)$, the vector space $\mathfrak{s}$ consists of $X \in M_{n+1}(E)$ such that

$$\overline{X} + X = 0.$$

Its analogous companion is the space $M_{n+1}(F)$ (or $\mathfrak{gl}_{n+1}(F)$) viewed as a subspace of $M_{n+1}(E)$, namely consisting of $X \in M_{n+1}(E)$ such that

$$\overline{X} = X.$$

For any number $\tau \in E$ such that $\overline{\tau} = -\tau \neq 0$, we have an isomorphism from $\mathfrak{s}(F)$ to $M_{n+1}(F)$ mapping X to $\tau^{-1}X$.

We have a canonical bi- E -linear pairing given by $(x, y) \mapsto \mathrm{tr}(xy) \in E$. We consider both $\mathfrak{s}$ and $\mathfrak{u}$ as F -vector subspaces of $M_{n+1}(E)$. The restrictions of the bilinear form on $M_{n+1}(E)$ to $\mathfrak{s}$ and $\mathfrak{u}$ take values in F and are non-degenerate. Moreover, if we change the Hermitian matrix defining $\mathfrak{u}$ to an equivalent one, the subspace $\mathfrak{u}$ changes to its conjugate by an element in $\mathrm{GL}_{n+1}(E)$. The additive character ψ and ψ_E determine a self-dual measure on $M_{n+1}(\mathbb{A}_E)$, $\mathfrak{u}(\mathbb{A})$, $\mathfrak{s}(\mathbb{A})$ and the local analogues. These measures can also be treated as the Tamagawa measure associated to differential forms. Let ω_0 be the differential form on $\mathfrak{u}$ that defines the self-dual measure. We use it to normalize the differential form that defines the measure on $U(\theta)(F_v)$. We consider the Cayley map

$$\mathfrak{c}(X) := (1 + X)(1 - X)^{-1}$$

defines a birational map between $\mathfrak{u}$ and $U(\theta)$, and it is defined at $X = 0$. We normalize the invariant differential form ω on $U(\theta)$ by requiring that the pull back $\mathfrak{c}^*\omega$ evaluating at 0 is the same as ω_0 evaluating at 0. It follows that, when v is non-archimedean, the Cayley map takes the restriction of the self-dual measure to a small neighborhood of 0 in $\mathfrak{u}$ to the restriction of the Tamagawa measure $|\omega|_v$ to a small neighborhood of 1 in $U(\theta)(F_v)$. In this way we define the Tamagawa measure on $U(\theta)(\mathbb{A})$, globally and locally, as follows:

$$dh = \prod_v L(1, \eta_v) |\omega|_v.$$

Finally, we note that under the Tamagawa measure, the volume of $[U(1)] = U(1)(F) \backslash U(1)(\mathbb{A})$ is given by

$$\text{vol}([U(1)]) = 2L(1, \eta).$$

3 Explicit local factorization of some periods

In this section we decompose several global linear forms on general linear groups into explicit products of local invariant linear forms. Nothing is original in this section but we need to determine all constants for the proof of the main results.

Invariant inner product. Let $\Pi = \Pi_n$ be a cuspidal automorphic representations of $H_n(\mathbb{A}_E)$. We recall some basic facts on the Whittaker model of $\Pi = \otimes_w \Pi_w$. We extend the additive character ψ_E to a character of $N_n(E)$ by

$$\psi_E(u) = \psi_E\left(\sum_{i=1}^{n-1} u_{i,i+1}\right), \quad u = (u_{i,j}) \in N_n(E).$$

Then the Fourier coefficient of ϕ is defined as

$$W_\phi(g) = \int_{N(E) \backslash N(\mathbb{A}_E)} \phi(ug) \overline{\psi_E}(u) du.$$

Then we have $W_\phi \in \mathcal{C}^\infty(N_n(\mathbb{A}_E) \backslash H_n(\mathbb{A}_E), \psi_E)$. The map $\phi \mapsto W_\phi$ realizes an equivariant embedding $\Pi \hookrightarrow \mathcal{C}^\infty(N_n(\mathbb{A}_E) \backslash H_n(\mathbb{A}_E), \psi_E)$ with image denoted by $\mathcal{W}(\Pi, \psi_E)$. For $\phi \in \Pi = \otimes_w \Pi_w$ We assume that W_ϕ is decomposable

$$W_\phi(g) = \prod_w W_w(g_w), \quad W_w \in \mathcal{C}^\infty(N_n(E_w) \backslash H_n(E_w), \psi_{E,w})$$

where w runs over all places of E .

We compare the unitary structure in the decomposition

$$\Pi \simeq \bigotimes_w \mathcal{W}(\Pi_w, \psi_{E,w}).$$

On Π we have the Petersson inner product, for $\phi, \phi' \in \Pi$:

$$\langle \phi, \phi' \rangle_{Pet} = \int_{Z_n(\mathbb{A}_E)H_n(E)\backslash H_n(\mathbb{A}_E)} \phi(g) \overline{\phi'(g)} dg.$$

On $\mathcal{W}(\Pi_w, \psi_{E,w})$ we have an invariant inner product defined by

$$\vartheta_w(W_w, W'_w) = \int_{N_{n-1}(E_w)\backslash H_{n-1}(E_w)} W_w \begin{pmatrix} h & \\ & 1 \end{pmatrix} \overline{W'_w} \begin{pmatrix} h & \\ & 1 \end{pmatrix} dh.$$

When Π_w and $\psi_{E,w}$ are unramified, the vectors $W = W'$ are fixed by $H_n(\mathcal{O}_{E,w})$ and normalized by $W(1) = 1$, we have

$$\vartheta_w = L(1, \Pi_w \times \tilde{\Pi}_w).$$

This can be deduced from [27] Therefore we define a normalized invariant inner product

$$\vartheta_w^{\natural}(W_w, W'_w) = \frac{\vartheta_w(W_w, W'_w)}{L(1, \Pi_w \times \tilde{\Pi}_w)}.$$

Then the product $\prod_w \vartheta_w^{\natural}$ converges and defines an invariant inner product on $\mathcal{W}(\Pi, \psi_E)$. It is a natural question to compare it with the Petersson inner product. We now recall a result of Jacquet–Shalika.

Proposition 3.1. *We have the following decomposition of the Petersson inner product in terms of the local inner product $\vartheta_w^{\natural}$:*

$$(3.1) \quad \langle \phi, \phi' \rangle_{Pet} = \frac{n \cdot \text{Res}_{s=1} L(s, \Pi \times \tilde{\Pi})}{\text{vol}(E^\times \backslash \mathbb{A}_E^1)} \prod_v \vartheta_v^{\natural}(W_w, W'_w),$$

where $W_\phi = \otimes_w W_w$ and $W_{\phi'} = \otimes_w W'_w$.

Proof. Up to a constant this is proved by [27, §4]. We thus recall their proof in order to determine this constant and this will also be used below to decompose the Flicker-Rallis period. We consider an Eisenstein series associated to a Schwartz function $\Phi \in \mathcal{C}_c^\infty(\mathbb{A}_E^n)$. let $e = (0, 0, \dots, 1) \in E^n$ and P is the mirabolic subgroup of H_n consisting of matrices whose last row is e . Set

$$f(g, s) = |g|^s \int_{\mathbb{A}_E^\times} \Phi(eag) |a|^{ns} d^\times a, \quad \text{Re}(s) \gg 0.$$

Consider the Epstein-Eisenstein series

$$(3.2) \quad E(g, \Phi, s) := \sum_{\gamma \in ZP(E) \backslash H_n(E)} f(\gamma g, s)$$

which is absolutely convergent when $\text{Re}(s) > 1$. Equivalently, we have

$$E(g, \Phi, s) = |g|^s \int_{E^\times \backslash \mathbb{A}_E^\times} \sum_{\xi \in E^n - \{0\}} \Phi(\xi ag) |a|^{ns} d^\times a.$$

(Note: this corresponds to the case $\eta = 1$ in [27, §4].) It has meromorphic continuation to $\mathbb{C}$ and has a simple pole at $s = 1$ with residue ([27, Lemma 4.2])

$$\frac{\text{vol}(E^\times \backslash \mathbb{A}_E^1)}{n} \widehat{\Phi}(0).$$

Note that the only non-explicit constant denoted by c in [27, Lemma 4.2] is the volume of $E^\times \backslash \mathbb{A}_E^1$. Now consider the zeta integral

$$I(s, \Phi, \phi, \phi') = \int_{Z(\mathbb{A}_E)H_n(E) \backslash H_n(\mathbb{A}_E)} E(g, \Phi, s) \phi(g) \overline{\phi'}(g) dg.$$

On one hand, it has a pole at $s = 1$ with residue

$$\frac{\text{vol}(E^\times \backslash \mathbb{A}_E^1)}{n} \widehat{\Phi}(0) \langle \phi, \phi' \rangle_{Pet}.$$

On the other hand, when $\text{Re}(s)$ is large, it is also equal to the following integral

$$(3.3) \quad \Psi(s, \Phi, W_\phi, W_{\phi'}) = \int_{N_n(\mathbb{A}_E) \backslash H_n(\mathbb{A}_E)} \Phi(eg) W_\phi(g) \overline{W_{\phi'}(g)} |\det(g)|^s dg$$

where, by [27, Prop. 2.3] for unramified data of (Φ_w, W_w, W'_w) and $\psi_{E,w}$ at a place w , normalized such that $W(1) = W(1) = \Phi_w(0) = 1$, we have

$$\Psi(s, \Phi, W_w, W'_w) = \text{vol}(K_w) L(s, \Pi_w \times \widetilde{\Pi}_w).$$

(Note: for our measure on $N_n(E_w)$, we have $\text{vol}(N_n(E_w) \cap K_w) = 1$.) From this we may deduce that

$$\Psi(s, \Phi, W_\phi, W_{\phi'}) = L(s, \Pi \times \widetilde{\Pi}) \prod_w \frac{\Psi(s, \Phi_w, W_{\phi,w}, W_{\phi',w})}{L(s, \Pi_w \times \widetilde{\Pi}_w)},$$

where the local factors are entire functions of s and for almost all w they are equal to one. Moreover, all local factors converge absolutely in the half plane $\text{Re}(s) > 1 - \epsilon$ for some $\epsilon > 0$ ([27]). From this we deduce that its residue at $s = 1$ is given by another formula

$$\text{Res}_{s=1} L(s, \Pi \times \widetilde{\Pi}) \prod_w \frac{\Psi(1, \Phi_w, W_{\phi,w}, W_{\phi',w})}{L(1, \Pi_w \times \widetilde{\Pi}_w)}.$$

From the two formulae of the residue, we first deduce that $\vartheta_w(W_{\phi,w}, W_{\phi',w})$ is $H_n(E_w)$ -invariant and second that

$$\Psi(1, \Phi_w, W_{\phi,w}, W_{\phi',w}) = \widehat{\Phi}_w(0) \vartheta_w(W_{\phi,w}, W_{\phi',w}).$$

We consider the action of $H_n(E)$ on the row vector space E^n from right multiplication. Then the stabilizer of e is the mirabolic subgroup P_n . Let $N_{n,1,+}(E_w)$ be the unipotent part of P_n and $N_{n,1,-}(E_w)$ the transpose of $N_{n,1,+}(E_w)$. We consider the open dense subset

$N_{n,1,+}H_{n-1}N_{n,1,-}Z_n = P_nN_{n,1,-}Z_n$. We may decompose the measure on H_n (or more precisely its restriction to the open subset)

$$dg = |\det(h)|^{-1}dn_+dhdn_-d^*a,$$

where

$$g = n_+hn_-a, \quad h \in H_{n-1}, n_{\pm} \in N_{n,1,\pm}, a \in Z_n.$$

Note also that the embedding $H_{n-1} \hookrightarrow P_n$ induces an isomorphism $N_n \backslash P_n \simeq N_{n-1} \backslash H_{n-1}$. For an integrable function f on $N_n \backslash H_n$, we may write

$$\begin{aligned} & \int_{N_n(E_w) \backslash H_n(E_w)} f(g)dg \\ &= \int_{Z_n N_{n,1,-}(E_w)} \left(\int_{N_{n-1}(E_w) \backslash H_{n-1}(E_w)} f(hn_-a) |\det(h)|^{-1}dh \right) dn_-d^*a \end{aligned}$$

We now apply this formula to the integral $\Psi(1, \Phi_w, W_{\phi,w}, W_{\phi',w})$. Clearly ϑ_w is $P(E_w)$ -invariant. Therefore we may write

$$\Psi(1, \Phi_w, W_{\phi,w}, W_{\phi',w}) = \int_{Z_n N_{n,1,-}(E_w)} \Phi(ean_-) \vartheta_w(\Pi_w(an_-)W_w, \Pi_w(an_-)W'_w) |a|^n d^*a dn_-.$$

For $X \in E_w^n$ (with last entry nonzero), let $n_-(X)$ be the element in $Z_n N_{n,1,-}(E_w)$ with last row equal to X . We consider

$$\Gamma(X) := \vartheta_w(\Pi_w(n_-(X))W_w, \Pi_w(n_-(X))W'_w),$$

whenever it is defined. A suitable substitution yields

$$\Psi(1, \Phi_w, W_{\phi,w}, W_{\phi',w}) = \int_{E_w^n} \Phi_w(X) \Gamma(X) dX.$$

Since by the other residue formula, we also know that this is equal to a constant multiple times $\widehat{\Phi}_w(0)$ times an invariant inner product on $\mathcal{W}(\Pi_w, \psi_{E,w})$, for all Φ_w and W_w, W'_w . We deduce that $\Gamma(X)$ is a constant function (whenever it is defined). Therefore $\Gamma(X) = \vartheta_w(W_w, W'_w)$ and ϑ_w is $H_n(E_w)$ -invariant. Moreover we now have

$$\Psi(1, \Phi_w, W_{\phi,w}, W_{\phi',w}) = \vartheta_w(W_w, W'_w) \int_{E_w^n} \Phi(X) dX = \vartheta_w(W_w, W'_w) \widehat{\Phi}_w(0).$$

This completes the proof. We also note that if we use the $H_n(E_w)$ -invariance of ϑ_w , which can be proved independently, then the proposition can be deduced immediately from the two residue formulae. From the $H_n E_w$ -invariance of ϑ_w and the equation (3.3), we may immediately deduce that $\vartheta_w = L(1, \Pi_w \times \widetilde{\Pi}_w)$ for unramified data, as promised earlier. $\square$

For later use, as we will be in the case of a quadratic extension of number fields, we will now consider $H_{n,E}$ as an algebraic group over the base field F . Therefore we rewrite the result as

$$(3.4) \quad \langle \phi, \phi' \rangle_{Pet} = \frac{n \cdot \text{Res}_{s=1} L(s, \Pi \times \tilde{\Pi})}{\text{vol}(E^\times \backslash \mathbb{A}_E^1)} \prod_v \vartheta_v^{\natural}(W_v, W'_v),$$

where $\vartheta_v = \prod_{w|v} \vartheta_w$ for all (one or two) places w above v .

Flicker-Rallis period. Now let E/F be a quadratic extension of number fields and $\Pi = \Pi_n$ a cuspidal automorphic representation of $H_n(\mathbb{A}_E)$. Assume that its central character $\omega_\Pi|_{\mathbb{A}^\times} = 1$. We would like to decompose the Flicker-Rallis period ([4],[9]) explicitly. It can be viewed as a twisted version of the Petersson inner product (it indeed gives the Petersson inner product if we allow $E = F \times F$ to be split globally). Therefore it is natural that the method is similar as well.

We first assume that n is odd. Then we have the global Flicker-Rallis period, an $H_n(\mathbb{A})$ -invariant linear form on Π :

$$(3.5) \quad \beta(\phi) = \beta_n(\phi) := \int_{Z_n(\mathbb{A})H_n(F) \backslash H_n(\mathbb{A})} \phi(h) dh, \quad \phi \in \Pi.$$

The global period β is related to the Asai L-function $L(s, \Pi, \text{As}^+)$ (for the definition of $\text{As}^\pm$, cf. [6]). We set

$$\tilde{\epsilon}_n = \text{diag}(\tau^{n-1}, \tau^{n-2}, \dots, 1) \in H_n(E)$$

and

$$\epsilon_{n-1} = \tau \cdot \text{diag}(\tau^{n-2}, \tau^{n-3}, \dots, 1) = \tau \tilde{\epsilon}_{n-1} \in H_{n-1}(E).$$

(Note: $\tau \in E^-$.) Indeed we may choose any $\tilde{\epsilon}_n = \text{diag}(a_1, \dots, a_{n-1}, a_n)$ such that $a_i/a_{i+1} \in E^-$ for $i = 1, \dots, n-1$ and $a_n = 1$. We again use the Whittaker model of Π . We will again consider $H_{n,E}$ as an algebraic group over F . In particular, we consider Π_v as a representation of $H_n(E_v)$ where $E_v = E \otimes_F F_v$ is a semisimple F_v algebra of rank two. For a place v of F , we define for $W_v \in \mathcal{W}(\Pi_v, \psi_{E_v})$

$$(3.6) \quad \beta_v(W_v) = \int_{N_{n-1}(F_v) \backslash H_{n-1}(F_v)} W_v \left(\begin{pmatrix} \epsilon_{n-1} h & \\ & 1 \end{pmatrix} \right) dh.$$

It depends on the choice of $\tau = \sqrt{\delta}$. For unramified data with normalization $W_w(e) = 1$, we have

$$\beta_v(W_v) = \text{vol}(K_n(\mathcal{O}_{F_v})) L(1, \Pi_v, \text{As}^+).$$

We thus define a normalized linear form

$$(3.7) \quad \beta_v^{\natural}(W_v) = \frac{\beta(W_v)}{L(1, \Pi_v, \text{As}^+)}.$$

Proposition 3.2. *We have an explicit decomposition*

$$(3.8) \quad \beta(\phi) = \frac{n \operatorname{Res}_{s=1} L(s, \Pi, \mathbb{A}s^+)}{\operatorname{vol}(F^\times \backslash \mathbb{A}^1)} \prod_v \beta_v^{\natural}(W_v),$$

where $W = W_\phi = \otimes_v W_v \in \mathcal{W}(\Pi, \psi_E)$.

Proof. Let $\Phi \in \mathcal{S}(\mathbb{A}^n)$ and we consider the same kind of Eisenstein series $E(g, \Phi, s)$ as in the previous paragraph, but now on $H_n(\mathbb{A})$. Then we define

$$I(s, \Phi, \phi) := \int_{Z(\mathbb{A})H_n(F) \backslash H_n(\mathbb{A})} E(g, \Phi, s) \phi(g) dg.$$

We then have ([4, pp.303])

$$(3.9) \quad I(s, \phi, \Phi) = \Psi(s, W_\phi, \Phi),$$

where

$$\Psi(s, W_\phi, \Phi) = \int_{N_n(\mathbb{A}) \backslash H_n(\mathbb{A})} W_\phi(\tilde{\epsilon}_n h) \Phi(eh) |h|^s dh.$$

(Note that we have different definition of the additive character ψ_E .) Indeed, we have

$$\begin{aligned} I(s, \Phi, \phi) &= \int_{P_n(F) \backslash H_n(\mathbb{A})} \Phi(eg) \phi(g) |g|^s dg \\ &= \int_{P_n(F) N_n(\mathbb{A}) \backslash H_n(\mathbb{A})} \Phi(eg) \left(\int_{N_n(F) \backslash N_n(\mathbb{A})} \phi(ng) dn \right) |g|^s dg. \end{aligned}$$

We have a Fourier expansion

$$\phi(g) = \sum_{\gamma \in N_n(E) \backslash P_n(E)} W_\phi(\gamma g).$$

Only those γ such that $\psi_E(\gamma n \gamma^{-1}) = 1$ for all $n \in N_n(\mathbb{A})$ contribute nontrivially. Therefore we may replace the sum by $\gamma \in \tilde{\epsilon}_n P_n(F)$:

$$\begin{aligned} I(s, \Phi, \phi) &= \int_{P_n(F) N_n(\mathbb{A}) \backslash H_n(\mathbb{A})} \Phi(eg) \left(\sum_{\gamma \in N_n(F) \backslash P_n(F)} W_\phi(\tilde{\epsilon}_n \gamma g) \right) |g|^s dg \\ &= \int_{N_n(\mathbb{A}) \backslash H_n(\mathbb{A})} \Phi(eg) W_\phi(\tilde{\epsilon}_n g) |g|^s dg \\ &= \Psi(s, W_\phi, \Phi). \end{aligned}$$

(Note that $\operatorname{vol}(N_n(F) \backslash N_n(\mathbb{A})) = 1$.) We define for each place v of F ,

$$\Psi(s, W_v, \Phi_v) = \int_{N_n(F_v) \backslash H_n(F_v)} W_v(\tilde{\epsilon}_n h) \Phi_v(eh) |h|^s dh.$$

For unramified data, we have

$$\Psi(s, W_v, \Phi_v) = \text{vol}(K_n(\mathcal{O}_{F_v}))L(s, \Pi_v, \text{As}^+).$$

And we have ([9, pp.185])

$$\Psi(1, W_v, \Phi_v) = \beta_v(W_v)\hat{\Phi}_v(0).$$

Alternatively we may prove this using (3.9), analogous to the proof of Prop. 3.1. Again, analogous to the proof of Prop. 3.1, we may take residue of (3.9) to obtain:

$$\frac{\text{vol}(F^\times \backslash \mathbb{A}^1)}{n} \hat{\Phi}(0) \beta(\phi) = \text{Res}_{s=1} L(s, \Pi, \text{As}^+) \hat{\Phi}(0) \prod_v \beta_v^\natural(W_v)$$

This completes the proof. $\square$

When n is even, we insert the character η in the definition of β

$$(3.10) \quad \beta(\phi) = \beta_n(\phi) := \int_{Z_n(\mathbb{A})H_n(F) \backslash H_n(\mathbb{A})} \phi(h) \eta(h) dh, \quad \phi \in \Pi.$$

The Asai L-function is then replaced by $L(s, \Pi, \text{As}^-)$, or we may write it as $L(s, \Pi, \text{As}^{(-1)^{n-1}})$. We also modify the definition

$$(3.11) \quad \beta_v(W_v) = \int_{N_{n-1}(F_v) \backslash H_{n-1}(F_v)} W_v \left(\begin{pmatrix} \epsilon_{n-1} h & \\ & 1 \end{pmatrix} \right) \eta_v(h) dh.$$

The same argument shows that (3.8) still holds.

Rankin-Selberg period. We now follow [25]. Let $\Pi = \Pi_n \otimes \Pi_{n+1}$ and Π_i a cuspidal automorphic representations of $H_i(\mathbb{A}_E)$, $i = n, n+1$. We define the global Rankin-Selberg period as

$$(3.12) \quad \lambda(\phi) = \int_{H_n(E) \backslash H_n(\mathbb{A}_E)} \phi(h) dh, \quad \phi \in \Pi,$$

where H_n embeds diagonally into $H_n \times H_{n+1}$. To decompose it, we need the Whittaker model $\mathcal{W}(\Pi_n, \bar{\psi}_E)$ ($\mathcal{W}(\Pi_{n+1}, \psi_E$, resp.) of Π_{n+1} with respect to the additive character $\bar{\psi}_E$ (ψ_E , resp.). We define a local Rankin-Selberg period on the local Whittaker model $W \in \mathcal{W}(\Pi_n, \bar{\psi}_E) \otimes \mathcal{W}(\Pi_{n+1}, \psi_E)$:

$$(3.13) \quad \lambda_w(W_w) = \int_{N_n(E_w) \backslash H_n(E_w)} W_w(h) dh,$$

and normalize it by

$$\lambda_w^\natural(W_w) = \frac{\lambda_w(W_w)}{L(1, \Pi_{n,w} \times \Pi_{n+1,w})}.$$

The integral is absolutely convergent for tempered representation. Then for unramified data we have $\lambda_w^{\natural}(W_w) = \text{vol}(K_w)$ ([25]). Finally, Jacquet, Piatetski-Shapiro and Shalika showed that

$$(3.14) \quad \lambda(\phi) = L\left(\frac{1}{2}, \Pi_n \times \Pi_{n+1}\right) \prod_w \lambda_w^{\natural}(W_w).$$

Decomposing a spherical character. We now denote

$$G' = \text{Res}_{E/F}(\text{GL}_n \times \text{GL}_{n+1})$$

viewed as an F -algebraic group. We consider its two subgroups: H'_1 is the diagonal embedding of $\text{Res}_{E/F}\text{GL}_n$ (where GL_n is embedded into GL_{n+1} by $g \mapsto \text{diag}[g, 1]$) and H'_2 is $\text{GL}_{n,F} \times \text{GL}_{n+1,F}$ embedded into G' in the obvious way.

Now we consider a cuspidal automorphic representation $\Pi = \Pi_n \otimes \Pi_{n+1}$ of $G'(\mathbb{A})$. We denote by $\mathcal{W}(\Pi, \psi)$ the Whittaker model $\mathcal{W}(\Pi_n, \bar{\psi}) \otimes \mathcal{W}(\Pi_{n+1}, \psi)$. Denote by $\beta = \beta_n \otimes \beta_{n+1}$ the (product of) Flicker–Rallis period on $\Pi = \Pi_n \otimes \Pi_{n+1}$. We define a *global spherical character*—a distribution on $H(\mathbb{A})$ —as follows. For $f' \in \mathcal{C}_c^\infty(G'(\mathbb{A}))$, define

$$(3.15) \quad I_\Pi(f') = \sum_{\phi} \frac{\lambda(\Pi(f')\phi)\overline{\beta(\phi)}}{\langle \phi, \phi \rangle_{\text{Pet}}},$$

where the sum runs over an *orthogonal* basis of Π . Equivalently,

$$I_\Pi(f') = \sum_{\phi} \lambda(\Pi(f')\phi)\overline{\beta(\phi)},$$

where the sum runs over an *orthogonal normal* basis of Π for the Petersson inner product. Note that the definition of $\Pi(f')$ involves a choice of the measure on $G'(\mathbb{A})$: we could choose any as long as then we use the same measure (quotient by the counting measure on $G'(F)$) to define the Petersson inner product.

By definition of $As^\pm$, we have for $i = n, n+1$:

$$L(s, \Pi_i \times \Pi_i^\sigma) = L(s, \Pi_i, As^+)L(s, \Pi_i, As^-).$$

Now assume that $\Pi = \pi_E$ is the base change of a cuspidal automorphic representation π of $G(\mathbb{A})$. By [6, Prop. 7.4] we also have

$$L(s, \Pi_i, As^{(-1)^i}) = L(s, \pi_i, ad).$$

Since Π is conjugate self-dual: $\tilde{\Pi} \simeq \Pi^\sigma$ where σ is the nontrivial element in $\text{Gal}(E/F)$, we deduce that

$$(3.16) \quad \frac{\text{Res}_{s=1} L(1, \Pi_i \times \tilde{\Pi}_i)}{\text{Res}_{s=1} L(s, \Pi_i, As^{(-1)^{i-1}})} = L(1, \Pi_i, As^{(-1)^i}) = L(1, \pi_i, ad),$$

where the function $L(s, \Pi_i, \text{As}^{(-1)^i})$ is regular and non-zero at $s = 1$ by a [35, Theorem 5.1] (cf. [21, Remark 1.4]).

We now define a (*normalized*) *local spherical character* for each place v of F :

$$(3.17) \quad I_{\Pi_v}^{\natural}(f'_v) = \sum_{W_v} \frac{\lambda_v^{\natural}(\Pi_v(f'_v)W_v) \overline{\beta_v^{\natural}(W_v)}}{\vartheta_v^{\natural}(W_v, W_v)}$$

where the sum runs over an orthogonal basis $W_v \in \mathcal{W}(\Pi_v, \psi_v)$ and λ_v, β_v are the local Rankin-Selberg periods via the Whittaker model.

We now summarize to arrive at an analogue of the decomposition in Conjecture [?]

Proposition 3.3. *Assume that the cuspidal automorphic representation Π of $G'(\mathbb{A})$ is the base change π_E of a cuspidal automorphic representation π of $G(\mathbb{A})$. Then we have*

$$(3.18) \quad I_{\Pi}(f') = L(1, \eta)^2 \frac{L(1/2, \Pi)}{L(1, \pi, \text{Ad})} \prod_v I_{\Pi_v}^{\natural}(f'_v).$$

Proof. Note that

$$\frac{\text{vol}(E^{\times} \backslash \mathbb{A}_E^1)}{\text{vol}(F^{\times} \backslash \mathbb{A}_F^1)} = L(1, \eta).$$

Then the result follows from Prop. 3.1, 3.2 and the relation (3.16). □

4 Relative trace formulae of Jacquet and Rallis

The construction of Jacquet and Rallis. We recall the Jacquet–Rallis relative trace formulae ([26]) and we refer to [50] for more details.

First we recall the construction of Jacquet–Rallis’ RTF in the unitary case. For $f \in \mathcal{C}_c^{\infty}(G(\mathbb{A}))$ we consider a kernel function

$$K_f(x, y) = \sum_{\gamma \in G(F)} f(x^{-1}\gamma y),$$

and a distribution

$$J(f) := \int_{H(F) \backslash H(\mathbb{A})} \int_{H(F) \backslash H(\mathbb{A})} K_f(x, y) dx dy.$$

A cuspidal automorphic representation π of $G(\mathbb{A})$ contributes to the spectral side

$$(4.1) \quad J_{\pi}(f) = \sum_{\phi} \mathcal{P}(\pi(f)\phi) \overline{\mathcal{P}(\phi)},$$

where the sum is now over an orthonormal basis of π with respect to the Petersson inner product:

$$\langle \phi, \phi' \rangle = \int_{G(F) \backslash G(\mathbb{A})} \phi(g) \overline{\phi'}(g) dg.$$

We call the distribution J_π a *global spherical character* on G (relative to $H \times H$). For a regular semisimple $\delta \in G(F)$ and $f \in \mathcal{C}_c^\infty(G(F))$, we define its orbital integral

$$O(\delta, f) = \int_{H(F) \times H(F)} f(x^{-1} \delta y) dx dy.$$

We now recall the RTF in the general linear case. Recall that $G' = \text{Res}_{E/F}(\text{GL}_n \times \text{GL}_{n+1})$ as an F -algebraic group. We consider its two subgroups: H'_1 is the diagonal embedding of $\text{Res}_{E/F} \text{GL}_n$ (where GL_n is embedded into GL_{n+1} by $g \mapsto \text{diag}[g, 1]$) and $H'_2 = \text{GL}_{n,F} \times \text{GL}_{n+1,F}$ embedded into G' in the obvious way. For $f' \in \mathcal{C}_c^\infty(G'(\mathbb{A}))$, we define a kernel function

$$K_{f'}(x, y) = \sum_{\gamma \in G'(F)} f'(x^{-1} \gamma y).$$

We then consider a distribution on $G'(\mathbb{A})$:

$$I(f') = \int_{H'_1(F) \backslash H'_1(\mathbb{A})} \int_{H'_2(F) \backslash H'_2(\mathbb{A})} K_{f'}(h_1, h_2) \eta(h_2) dh_1 dh_2.$$

A cuspidal automorphic representation Π of $G'(\mathbb{A})$ contributes to the spectral side

$$(4.2) \quad I_\Pi(f') = \sum_{\phi} \mathcal{P}_1(\Pi(f')\phi) \overline{\mathcal{P}_2(\phi)},$$

where $\mathcal{P}_1 = \lambda$, $\mathcal{P}_2 = \beta_n \otimes \beta_{n+1}$ correspond the global Rankin-Selberg and Flicker-Rallis periods respectively, and the sum is now over an orthonormal basis of Π (with respect to the Petersson inner product). This is of course the same as the spherical character we defined earlier in (3.17).

We also recall the definition of orbital integrals for an $H'_1 \times H'_2$ -regular semisimple (cf. [49, §2])—for simplicity, *regular semisimple*:

$$(4.3) \quad O(\gamma, f') := \int_{H'_1(F)} \int_{H'_2(F)} f'(h_1^{-1} \gamma h_2) \eta(h_2) dh_1 dh_2.$$

Denote by $(H_1(F) \backslash G'(F) / H_2(F))_{rs}$ the set of regular semisimple $(H'_1 \times H'_2)(F)$ -orbits in $G'(F)$ and $(H(F) \backslash G(F) / H(F))_{rs}$ the set of regular semisimple $(H \times H)(F)$ -orbits in $G(F)$. To show the dependence on the Hermitian spaces W, V , we will write $G_{W,V}$ for G and H_W for H . Then there is a natural bijection

$$(H_1(F) \backslash G'(F) / H_2(F))_{rs} \simeq \coprod_{W \subset V} (H_W(F) \backslash G_{W,V}(F) / H_W(F))_{rs},$$

where on the RHS the disjoint union runs over all pair $W \subset V$ of Hermitian spaces of dimension n and $n+1$ modulo the equivalence relation: (W, V) is equivalent to (W', V') if there a constant $\kappa \in F^\times$ such that $\kappa W \simeq W'$ and $\kappa V \simeq V'$ (here κW means that we multiply the Hermitian form by the constant κ). Without loss of generality, we may and will assume that $V = W \oplus Ee$ is an orthogonal sum of W and a one-dimensional Hermitian space Ee with a norm one vector $\langle e, e \rangle = 1$. Moreover, the same holds if we replace F by F_v for every place v of F (for the proof, see [49, §2]).

There is an explicit *transfer factor* $\{\Omega_v\}_v$ defined in [50, §2.4] with the following property on the regular semisimple locus of $G'(F_v)$, indexed by all places v of F , such that

- If $\gamma \in G'(F)$ is regular semisimple, then we have a product formula $\prod_v \Omega_v(\gamma) = 1$.
- For any $h_i \in H'_i(F_v)$ and $\gamma \in G'(F_v)$, we have $\Omega(h_1 \gamma h_2) = \eta(h_2) \Omega_v(\gamma)$.

We may construct it as follows. Let S_{n+1} is the subvariety of $Res_{E/F} GL_{n+1}$ defined by the equation $s\bar{s} = 1$. By Hilbert Satz-90, we have an isomorphism of two affine varieties

$$Res_{E/F} GL_{n+1} / GL_{n+1, F} \simeq S_{n+1},$$

induced by the morphism between F -varieties:

$$\begin{aligned} \nu : Res_{E/F} GL_{n+1} &\rightarrow S_{n+1} \\ g &\mapsto g\bar{g}^{-1}, \end{aligned}$$

and in the level of F -points:

$$GL_{n+1}(E) / GL_{n+1}(F) \simeq S_{n+1}(F).$$

Fix an auxiliary character throughout this paper:

$$(4.4) \quad \eta' : E^\times \backslash \mathbb{A}_E^\times \rightarrow \mathbb{C}^\times$$

(not necessarily quadratic) such that its restriction $\eta'|_{\mathbb{A}^\times} = \eta$. Write $\gamma = (\gamma_1, \gamma_2) \in G'(F_v)$ and $s = \nu(\gamma_1 \gamma_2^{-1})$. If n is odd, we set:

$$(4.5) \quad \Omega_v(\gamma) := \eta'_v(\det(\gamma_1 \gamma_2^{-1})) \eta'_v(\det(s)^{-(n+1)/2} \det(es, \dots, es^n)),$$

and if n is even, we set:

$$(4.6) \quad \Omega_v(\gamma) := \eta'_v(\det(s)^{-n/2} \det(es, \dots, es^n)).$$

For a place v of F , we say that the function $f' \in \mathcal{C}_c^\infty(G'(F_v))$ and the tuple $(f_{W,V})_{(W,V)}, f_{W,V} \in \mathcal{C}_c^\infty(G_{W,V}(F_v))$, indexed by the set of all equivalence classes of Hermitian pair (W, V) over $E_v = E \otimes F_v$, are *smooth transfer of each other* or *match* if

$$\Omega_v(\gamma) O(\gamma, f') = O(\delta, f_{W,V}),$$

whenever a regular semisimple $\gamma \in G'(F_v)$ matches $\delta \in G_{W,V}(F_v)$. One of the main local results in [50] is the existence of smooth transfer at non-archimedean non-split places (cf. [50, Theorem 2.6]) and arbitrary split places (cf. [50, Prop. 2.5]). In this paper, we usually need to consider a fixed pair (W, V) and we say that f' and $f_{W,V} \in \mathcal{C}_c^\infty(G_{W,V}(F_v))$ match if there exists some $f_{W',V'}$ for all equivalence classes $(W', V') \neq (W, V)$ such that f' match the completed tuple $f_{W,V}, f_{W',V'}$. Moreover, by [44], the fundamental lemma holds when the characteristic of the residue field at v is sufficiently large.

We need some simplification ([50]). Identify $H'_1 \backslash G'$ with $\text{Res}_{E/F} \text{GL}_{n+1}$. Now we write F for F_v for a fixed place v . We may integrate f' over $H'_1(F)$ to get a function on $\text{Res}_{E/F} \text{GL}_{n+1}(F)$:

$$\bar{f}'(x) := \int_{H'_1(F)} f'((x, 1)h_1) dh_1, \quad x \in \text{Res}_{E/F} \text{GL}_{n+1}(F).$$

Now assuming that n is odd, so the character η on H'_2 is indeed only nontrivial on the component $\text{GL}_{n+1,F}$. Then we may introduce a function on $S_{n+1}(F)$ as follows: when $\nu(x) = s \in S_{n+1}(F)$, we define

$$\tilde{f}'(x) := \int_{\text{GL}_{n+1}(F)} \bar{f}'(xg) \eta'(xg) dg.$$

Then $\tilde{f}' \in \mathcal{C}_c^\infty(S_{n+1}(F))$ and all functions in $\mathcal{C}_c^\infty(S_{n+1}(F))$ arise this way. Now it is easy to see that for $\gamma = (\gamma_1, \gamma_2)$:

$$(4.7) \quad O(\gamma, f') = \eta'(\det(\gamma_1 \gamma_2^{-1})) \int_{\text{GL}_n(F)} \tilde{f}'(h^{-1}sh) \eta(h) dh, \quad s = \nu(\gamma_1 \gamma_2^{-1}).$$

If n is even, we simply define in the above

$$\tilde{f}'(x) := \int_{\text{GL}_{n+1}(F)} \bar{f}'(xg) dg,$$

and then

$$(4.8) \quad O(\gamma, f') = \int_{\text{GL}_n(F)} \tilde{f}'_v(h^{-1}sh) \eta(h) dh, \quad s = \nu(\gamma_1 \gamma_2^{-1}).$$

Up to a sign, the integral on the RHS depends only on the orbit of s under the conjugation by $\text{GL}_n(F)$. Moreover, in either case we have

$$\Omega(\gamma) O(\gamma, f') = \Omega(s) O(s, \tilde{f}'), \quad s = \nu(\gamma_1 \gamma_2^{-1}).$$

A trace formula identity. We are lead to a comparison of the two RTFs and the two spherical characters I_Π and J_π when Π is the base change of π .

Conjecture 4.1. *Let π be tempered automorphic cuspidal representation on $G(\mathbb{A})$ that admits invariant linear functional:*

$$\mathrm{Hom}_{H(\mathbb{A})}(\pi, \mathbb{C}) \neq 0.$$

Let π_E be the base change of π . If π_E is cuspidal, then we have

$$2^{-2}L(1, \eta)^{-2}I_{\pi_E}(f') = J_{\pi}(f),$$

for matching f' and f .

Remark 5. If π_E is not cuspidal, then π is endoscopic and in the above equality the constant 2^2 should be replaced by $|S_{\pi}|$ appeared in the Conjecture 1.1. All of these should follow from the full spectral decomposition of the Jacquet-Rallis relative trace formulae.

Theorem 4.2. *The conjecture 4.1 holds if we assume that*

- *For at least one split place v , π_v is supercuspidal.*
- *If $v|\infty$, either v is split, or the group $G(F_v)$ is compact and f'_v matches the tuple $(f_v, 0, 0, \dots, 0)$ (with zero functions on every unitary group other than $G(F_v)$) satisfying that: the function f'_v is supported in the (open) subset of regular semisimple locus matching elements in the compact unitary group $G(F_v)$, and the function f_v is supported in the regular semisimple locus of $G(F_v)$.*

Proof. We would like to apply the result from [50]. But we need to compare the difference on the normalization of Petersson inner product in the unitary case (caused by the presence of center). There implicitly we use a different Petersson inner product

$$\langle \phi, \phi' \rangle' = \int_{Z(\mathbb{A})G(F)\backslash G(\mathbb{A})} \phi(g)\overline{\phi'}(g)dg = \mathrm{vol}(Z(F)\backslash Z(\mathbb{A}))^{-1}\langle \phi, \phi' \rangle.$$

Note that the center Z of G is isomorphic to $U(1) \times U(1)$. Hence its Tamagawa number is

$$\mathrm{vol}(Z(F)\backslash Z(\mathbb{A})) = (2L(1, \eta))^2.$$

Now taking into this correction, we apply the trace formula identity [50, Prop. 2.11] to matching f' and a tuple (f_W) indexed by equivalence classes of (W, V) :

$$I_{\pi_E}(f') = (2L(1, \eta))^2 \sum_W \sum_{\pi_W} J_{\pi_W}(f_W),$$

where the sum is over all equivalence classes of (W, V) and all π_W nearly equivalence to π . We assume that for the (W, V) we start with, $f_W = f$ is also the function in the assumption of the theorem. Note that by our assumption on L-packet, the multiplicity one holds, namely, each π_W are non-isomorphic. Moreover, if an archimedean place v is split, $\pi_{W,v}$ is isomorphic to π_v (there is only one member in the L-packet of the general linear group). If an archimedean

place v is nonsplit, then the compactness of $G(F_v)$ and the assumption on the matching f'_v and the tuple $(f_v, 0, 0, \dots, 0)$ again implies that $\pi_{W,v}$ is isomorphic to π_v (we are using the fact that for the compact unitary group, the Vogan L-packet of π_v contains only one member). If v is a p -adic place, the result on local Gan-Gross-Prasad in [2] assures that there is at most one member in the same Vogan L-packet of π_v admitting invariant linear functional. By our assumption $\text{Hom}_{H(F_v)}(\pi_v, \mathbb{C}) \neq 0$. Hence the sum reduces to one term

$$I_{\pi_E}(f') = (2L(1, \eta))^2 J_{\pi}(f).$$

□

Reduction to a local question. Our the main ingredient is an identity between the two local distributions. Let $d_n = n(n-1)(n-2)/6$ be the integer defined by

$$\tau^{d_n} = \delta_{n-1}(\epsilon_{n-1}) = \det(\text{Ad}(\epsilon_{n-1}) : N_{n-1}(E)).$$

Conjecture 4.3. *Let π_v be an irreducible tempered unitary representation of $G(F_v)$ with $\alpha_v \neq 0$. Assume that the base change Π_v is irreducible generic unitary (so that I_{Π_v} is well-defined). If the functions $f_v \in \mathcal{C}_c^\infty(G(F_v))$ and $f'_v \in \mathcal{C}_c^\infty(G'(F_v))$ match, then we have*

$$I_{\Pi_v}(f'_v) = \kappa_v L(1, \eta_v)^{-1} J_{\pi_v}(f_v),$$

where the constant κ_v is given by

$$\kappa_v = \kappa_v(\eta', \tau, n, \psi) = |\tau|_{E,v}^{(d_n + d_{n+1})/2} (\epsilon(1/2, \eta_v, \psi_v) / \eta'(\tau))^{n(n+1)/2} \eta_v(\text{disc}(W)) \omega_{\Pi_n}(\tau).$$

Here ω_{Π_n} is the central character of Π_n , and $\text{disc}(W) \in F^\times / \text{NE}^\times$ is the discriminant of the Hermitian space W .

Proposition 4.4. *The two conjectures 4.1 and 4.3 together imply Conjecture 1.4 and 1.1 when π_E is cuspidal.*

Proof. By Conjecture 4.1 and Prop. 3.3 we have

$$J_{\pi}(f) = 2^{-2} L(1, \eta)^{-2} I_{\pi_E}(f') = 2^{-2} \frac{L(1/2, \pi_E)}{L(1, \pi, \text{Ad})} \prod_v I_{\Pi_v}^{\natural}(f'_v).$$

Conjecture 4.3 is equivalent to the identity between the normalized distributions

$$I_{\Pi_v}^{\natural}(f'_v) = \kappa_v L(1, \eta_v)^{-1} \Delta_{n+1,v} J_{\pi_v}^{\natural}(f_v).$$

Since

$$\prod_v \epsilon(1/2, \eta_v, \psi_v) = \epsilon(1/2, \eta) = 1,$$

we have

$$\prod_v \kappa_v = 1.$$

Note that the product $\prod_v L(1, \eta_v)^{-1} \Delta_{n+1, v}$ converges absolutely to $L(1, \eta)^{-1} \Delta_{n+1}$. We thus obtain

$$J_\pi(f) = 2^{-2} L(1, \eta)^{-1} \Delta_{n+1} \frac{L(1/2, \pi_E)}{L(1, \pi, \text{Ad})} \prod_v J_{\pi_v}^\natural(f_v).$$

Note that the global measure on H and G in Introduction are normalized by $L(1, \eta)^{-1}$ and $L(1, \eta)^{-2}$ respectively. Correction of measures yields Conjecture 1.4 and therefore Conjecture 1.1. $\square$

We have the following evidence of Conjecture 4.3.

Theorem 4.5. *Let v be a non-archimedean place and π_v as in Conjecture 4.3. Then the local conjecture 4.3 holds if the place v is split in E/F . Assuming that the ratio $I_{\Pi_v}(f'_v)/J_{\pi_v}(f_v)$ is a constant, then the local conjecture 4.3 holds under any one of the following conditions:*

- *The representation π_v is unramified, or*
- *The place v is non-split and $H(F_v)$ is compact, or*
- *The place v is non-split and the representation π_v is supercuspidal.*

Below we first prove the local conjecture when π_v is unramified or v is split and non-archimedean. We postpone the proof of the case v non-split to the last part of §9. When the place v is archimedean and split in E/F , we only have an equality in terms of α'_v instead of α_v . That's why in the main Theorem 1.2 we only use α'_v . We will prove Theorem 1.2 after Lemma 4.7.

The case of π_v unramified. Then we may assume that

1. The quadratic extension E/F is unramified at v .
2. The number τ is a v -adic unit.
3. The character ψ is unramified and hence so is ψ_E .

Indeed, it is easy to see how I_{Π_v} depends on τ . And if we twist ψ by $a \in F_v^\times$, it amounts to change τ by $a\tau$.

We need to utilize the fundamental lemma proved by Yun in the equal characteristic case and extended by J. Gordon's to the mixed character case (when the residue characteristic is large). We have a matching pair ⁵

$$f_v = \frac{1}{\text{vol}(H(\mathcal{O}_v))^2} 1_{G(\mathcal{O}_v)}, \quad f'_v = \frac{1}{\text{vol}(H'_1(\mathcal{O}_v))\text{vol}(H'_2(\mathcal{O}_v))} 1_{G'(\mathcal{O}_v)}.$$

⁵Note that the measures in the fundamental lemma proved in [44] are different from ours.

Let Π_v be the base change of π_v . Let $W_0 \in \mathcal{W}(\Pi_v, \psi_E)$ be the unique spherical element normalized such that $W_0(1) = 1$. Then we have

$$\Pi_v(f')W_0 = \frac{\text{vol}(G'(\mathcal{O}_v))}{\text{vol}(H'_1(\mathcal{O}_v))\text{vol}(H'_2(\mathcal{O}_v))}W_0.$$

and

$$I_{\Pi_v}(f'_v) = \frac{\lambda(\Pi_v(f'_v)W_0)\bar{\beta}(W_0)}{\vartheta_v(W_0, W_0)} = \frac{\text{vol}(G'(\mathcal{O}_v))}{\text{vol}(H'_1(\mathcal{O}_v))\text{vol}(H'_2(\mathcal{O}_v))} \frac{\lambda(W_0)\bar{\beta}(W_0)}{\vartheta_v(W_0, W_0)}.$$

Note that

$$\lambda(W_0) = L(1/2, \Pi_v) \cdot \text{vol}(H'_1(\mathcal{O}_v))$$

and

$$\frac{\bar{\beta}(W_0)}{\vartheta_v(W_0, W_0)} = L(1, \pi_v, Ad)^{-1} \frac{\text{vol}(H'_2(\mathcal{O}_v))}{\text{vol}(H_n(\mathcal{O}_{E,v}))\text{vol}(H_{n+1}(\mathcal{O}_{E,v}))}.$$

We obtain

$$I_{\Pi_v}(f'_v) = \frac{L(1/2, \Pi_v)}{L(1, \pi_v, Ad)} \cdot \frac{\text{vol}(G'(\mathcal{O}_v))}{\text{vol}(H'_1(\mathcal{O}_v))\text{vol}(H'_2(\mathcal{O}_v))} \cdot \frac{\text{vol}(H'_1(\mathcal{O}_v))\text{vol}(H'_2(\mathcal{O}_v))}{\text{vol}(H_n(\mathcal{O}_{E,v}))\text{vol}(H_{n+1}(\mathcal{O}_{E,v}))}.$$

In summary we have in the general linear case

$$I_{\Pi_v}(f'_v) = \frac{L(1/2, \Pi_v)}{L(1, \pi_v, Ad)}.$$

In the unitary case, we let $\phi_0 \in \pi_v^{K_v}$ normalized by $\|\phi_0\| = 1$:

$$\pi_v(f_v)\phi_0 = \frac{\text{vol}(G(\mathcal{O}_v))}{\text{vol}(H(\mathcal{O}_v))^2}\phi_0.$$

Therefore we have

$$J_{\pi_v}(f_v) = \alpha_v(\pi_v(f_v)\phi_0, \phi_0) = \frac{\text{vol}(G(\mathcal{O}_v))}{\text{vol}(H(\mathcal{O}_v))^2}\alpha_v(\phi_0, \phi_0).$$

By the unramified computation in [21]:

$$\alpha_v(\phi_0, \phi_0) = \text{vol}(H(\mathcal{O}_v))\Delta_{n+1,v} \frac{L(1/2, \Pi_v)}{L(1, \pi_v, Ad)}.$$

We thus obtain

$$J_{\pi_v}(f_v) = \frac{\text{vol}(G(\mathcal{O}_v))}{\text{vol}(H(\mathcal{O}_v))}\Delta_{n+1,v} \frac{L(1/2, \Pi_v)}{L(1, \pi_v, Ad)}.$$

Note that Δ_{n+1}^{-1} is the volume of $K_{n+1,v}$ for the local Tamagawa measure

$$\text{vol}(K_{n+1,v}) = L(1, \eta)\Delta_{n+1,v}^{-1}.$$

Therefore we obtain that

$$J_{\pi}(f) = L(1, \eta_v) \frac{L(1/2, \Pi_v)}{L(1, \pi_v, Ad)}.$$

We now complete the proof that

$$I_{\Pi_v}(f'_v) = L(1, \eta_v)^{-1}J_{\pi_v}(f_v).$$

Change of measures. *From now on, all measures will be the unnormalized (namely, without the convergence factor $\zeta_v(1), L(1, \eta_v)$ etc.) Tamagawa measures with the natural invariant differential forms on the general linear groups, their subgroups, and Lie algebras.*

Lemma 4.6. *When using the unnormalized measures, the identity in Conjecture in 4.3 becomes*

$$(4.9) \quad I_{\Pi_v}(f'_v) = \kappa_v J_{\pi_v}(f_v),$$

for matching functions f_v and f'_v (also under the unnormalized measures).

Proof. We then see that the old distribution I_{Π_v} is the new one times

$$\zeta_{E,v}(1)^2 \frac{\zeta_{E,v}(1) \zeta_{F,v}(1)^2}{\zeta_{E,v}(1)^2},$$

where the first term comes from the measure on G' involving the definition of $\Pi_v(f'_v)$, and the fraction comes from the measures in λ_v, β_v and ϑ_v . Similarly, the old distribution J_{π_v} is the new one times

$$L(1, \eta_v)^2 \cdot L(1, \eta_v),$$

where the first term comes from the measure on G involving the definition of $\pi_v(f_v)$, and the second from the measure on $H(F_v)$ in the definition of α_v . Moreover, the change of measures on $H'_1(F_v), H'_2(F_v)$ and $H(F_v)$ also change the requirement of smooth matching: if f_v and f'_v match for the normalized measures, then $\zeta_{E,v}(1) \zeta_{F,v}(1)^2 f_v$ and $L(1, \eta_v)^2 f'_v$ match for the unnormalized measures. Therefore, when using the unnormalized measures, the identity in Conjecture in 4.3 becomes the asserted one. $\square$

The case of a split place v . Assume that $F = F_v$ is split. Let $\pi = \pi_n \otimes \pi_{n+1}$ be an irreducible unitary generic representations of $G(F)$. We may identify $H_n(E)$ with $\mathrm{GL}_n(F) \times \mathrm{GL}_n(F)$ and identify $U(W)(F_v)$ with a subgroup consisting of elements of the form $(g, {}^t g^{-1})$, $g \in \mathrm{GL}_n(F)$ and ${}^t g$ is the transpose of g . Let p_1, p_2 be the two isomorphisms between $U(W)(F)$ with $\mathrm{GL}_n(F)$ induced by the two projections from $\mathrm{GL}_n(F) \times \mathrm{GL}_n(F)$ to $\mathrm{GL}_n(F)$. If π_n is an irreducible generic representation of $U(W)(F_v)$, the representation $\Pi_n = BC(\pi_n)$ can be identified with $p_1^* \pi_v \otimes p_2^* \pi_v$ of $H_n(E)$ where $p_i^* \pi_v$ is a representation of $\mathrm{GL}_n(F)$ obtained by the isomorphism p_i . For simplicity, we will write $\Pi_n = \pi_n \otimes \tilde{\pi}_n$. Similarly for π_{n+1}, Π_{n+1} . We fix a Whittaker model $\mathcal{W}(\pi_i), i = n, n+1$ using the additive character ψ at v . Recall that in the Introduction we have defined an alternative element $\alpha' \in \mathrm{Hom}(\pi \otimes \tilde{\pi}, \mathbb{C})$ by

$$\alpha'(W, W') = \lambda(W) \overline{\lambda(W')}, W, W' \in \mathcal{W}(\pi).$$

Again we have identified $\tilde{\pi}$ with $\bar{\pi}$. Moreover, we may require that the invariant inner product on π is the one defined by ϑ (cf. §3).

Lemma 4.7. *Let $f' = f_1 \otimes f_2 \in \mathcal{C}_c^\infty(G'(F))$ and $f = f_1 * f_2^* \in \mathcal{C}_c^\infty(G(F))$ ($f_2^*(x) = \overline{f_2}(x^{-1})$) matching f' . Then we have*

$$I_{\Pi_v}(f') = |\tau|_{E,v}^{(d_n+d_{n+1})/2} J'_\pi(f).$$

Moreover, Theorem 4.5 holds for split v .

Proof. We identify Π_n with $\pi_n \otimes \tilde{\pi}_n = \pi_n \otimes \bar{\pi}_n$. Then we have for $W, W' \in \mathcal{W}(\pi_n)$:

$$\beta_n(W \otimes \overline{W'}) = \int_{N_{n-1}(F) \backslash H_{n-1}(F)} W \begin{pmatrix} \epsilon_{n-1}h & \\ & 1 \end{pmatrix} \overline{W'} \begin{pmatrix} \epsilon_{n-1}h & \\ & 1 \end{pmatrix} dh.$$

This yields

$$\beta_n(W \otimes \overline{W'}) = |\tau|^{d_n/2} \vartheta_n(W, W').$$

Similarly for β_{n+1} . Then the desired equality follows by the definition of I_Π in terms of the linear functional λ, β and ϑ (note that $\delta = \tau^2$ is indeed a square in F). Now Theorem 4.5 follows from $\omega_{\Pi_n}(\tau) = \omega_{\pi_n} \cdot \omega_{\tilde{\pi}_n}(\tau) = 1$ since the central character $\omega_{\tilde{\pi}_n} = \omega_{\pi_n}^{-1}$. $\square$

we now give the proof of the first part of Theorem 1.2.

Proof of Theorem 1.2: the case (1). If for some place v the linear form α_v vanishes, then both sides vanish and Theorem 1.2 is trivial. Then by the multiplicity one result in [2], there does not exist non-zero $H(\mathbb{A})$ -invariant linear functional on any other member different from π in the Vogan L-packet of π . It follows from [50, Theorem 1.1] that if the linear functional $\mathcal{P}$ vanishes on π , the value $L(1/2, \pi_E)$ must vanish, or equivalently $\mathcal{L}(1/2, \pi_E) = 0$ (since the adjoint L-function does not vanish at $s = 1$). Hence we may further assume that the linear functional $\mathcal{P}$ does not vanish on π and $\mathcal{L}(1/2, \pi) \neq 0$. Under the condition of Theorem 1.2, Conjecture 4.1 holds by Theorem 4.2:

$$2^{-2} L(1, \eta)^{-2} I_{\pi_E}(f') = J_\pi(f).$$

It follows that the both sides are non-zero distribution. A prior we know that

$$J_\pi(f) = \mathcal{C}_\pi \prod_v J_{\pi_v}^\natural(f_v).$$

Therefore for every v , $I_{\Pi_v}(f'_v)$ differs from $J_{\pi_v}(f_v)$ by a non-zero constant independent of the matching functions f_v and f'_v . It follows from Theorem 4.5 that for all non-archimedean v , the identity in Conjecture 4.3 holds for $v|\infty$ when we replace J_{π_v} (using α_v) by J'_{π_v} (using α'_v). Since all $v|\infty$ are split in E/F , by Lemma 4.7, the identity also holds for $v|\infty$. Now Theorem 1.2 follows from the proof of Prop. 4.4. $\square$

Now we assume that F is *non-archimedean*. We would like to identify the distribution J'_π (defined using α') with J_π (defined using α). Note that in the definition of α , the inner product is now defined by ϑ :

$$\alpha(W, W') = \int_{H(F)} \langle \pi(h)W, W' \rangle dh, \quad \langle W, W' \rangle = \vartheta(W, W').$$

And the measure on $H(F) \simeq \mathrm{GL}_n(F)$ is the unnormalized Tamagawa measure. Let $c \in \mathbb{R}_+$ and let N^c be the subgroup of $N_n(F)$ consisting of u such that $|u_{i,i+1}| < c$, $i = 1, 2, \dots, n-1$.

Lemma 4.8. *Assume that π_n is tempered. For $W, W' \in \mathcal{W}(\pi_n)$, then the integral*

$$\int_{N^c(F)} \langle \pi_n(u)W, W' \rangle \psi(u^{-1}) du$$

stabilizes as $c \rightarrow \infty$. We define

$$\xi(W, W') := \int_{N_n(F)}^{reg} \langle \pi_n(u)W, W' \rangle \psi(u^{-1}) du$$

being the limit as $c \rightarrow \infty$. Then we have

$$\xi(W, W') = W(1) \overline{W'}(1).$$

Proof. The LHS does not vanish identically by Waldspurger [40, §3.5]. Both sides define non-zero elements in

$$\mathrm{Hom}_{N_n(F)}(\mathcal{W}_{\pi_n}, \psi) \otimes \mathrm{Hom}_{N_n(F)}(\mathcal{W}_{\overline{\pi_n}}, \overline{\psi}).$$

By uniqueness of Whittaker functional (up to scalar), the lemma holds up to a unique non-zero constant multiple. Then the ratio is evaluated by Lapid-Mao. $\square$

Proposition 4.9. *Assume that $\pi = \pi_n \otimes \pi_{n+1}$ is tempered. Then we have, for all $W, W' \in \mathcal{W}(\pi)$:*

$$\alpha(W, W') = \lambda(W) \overline{\lambda(W')}.$$

Namely, $\alpha = \alpha'$ as non-zero elements in $\mathrm{Hom}(\pi \otimes \tilde{\pi}, \mathbb{C})$.

Remark 6. This should also be true for F archimedean. But some convergence issue has not been solved. In practical application, it seems that the linear form α' is more suitable for computation than α .

Proof. The RHS does not vanish by the non-vanishing of local Rankin-Selberg integral ([25]). By the multiplicity one theorem, the LHS is a constant multiple of RHS for all W, W' . Hence it suffices to prove the identity for some choice of W, W' . Let $W = W_n \otimes W_{n+1}$, $W' = W'_n \otimes W'_{n+1}$. We choose W_{n+1}, W'_{n+1} such that their restriction to $H_n(F)$ has support compact modulo $N_n(F)$. More specifically, we fix an open (relative) compact subset $\Xi \subset A_n(F)N_{n-}(F)$ we may choose W_{n+1}, W'_{n+1} such that their restriction to $H_n(F)$ has support inside $N_n(F)\Xi$. Then we have, for $g \in H_n(F)$,

$$\langle \pi_{n+1}(g)W_{n+1}, W'_{n+1} \rangle = \int_{\Xi} W_{n+1}(xg) \overline{W'_{n+1}}(x) dx$$

for a suitable measure on Ξ . Let $c \in \mathbb{R}_+$ and consider the subgroup N^c of $N_n(F)$. Let us consider the integral

$$I_c(W, W') := \int_{\Xi} \int_{\Xi} \int_{N^c} \Phi_{W_n, W'_n}(x^{-1}ug) \psi(u) W_{n+1}(g) \overline{W'_{n+1}}(x) du dx dg.$$

This is equal to

$$I_c(W, W') := \int_{\Xi} \int_{\Xi} \int_{N^c} \Phi_{\pi_n(g)W_n, \pi_n(x)W'_n}(u) \psi(u) W_{n+1}(g) \overline{W'_{n+1}}(x) du dx dg.$$

By [40, §3] for non-archimedean case, the integral

$$(4.10) \quad \int_{N^c} \Phi_{\pi_n(g)W_n, \pi_n(x)W'_n}(u) \psi(u) du$$

converges absolutely. Its proof also shows that the integral defines a continuous function of both x and g . In particular, the integral I_c converges absolutely. Moreover, the limit of $I_c(W, W')$ as c goes to ∞ exists and is equal to

$$\int_{\Xi} \int_{\Xi} \pi_n(g)W_n(1) \pi_n(x)W'_n(1) W_{n+1}(g) \overline{W'_{n+1}}(x) dx dg.$$

Indeed, when F is non-archimedean, for each (x, g) , the integral in 4.10 stabilizes when c is large enough. But since x, g vary in a compact set Ξ , this constant c can be chosen uniformly. Therefore the limit $\lim_{c \rightarrow \infty} I_c(W, W')$ is equal to

$$\int_{\Xi} \int_{\Xi} \xi(\pi(g)W_n, \pi(x)W'_n) W_{n+1}(g) \overline{W'_{n+1}}(x) dx dg.$$

By Lemma 4.8 we have

$$\int_{\Xi} \int_{\Xi} W_n(g) \overline{W'_n}(x) W_{n+1}(g) \overline{W'_{n+1}}(x) dx dg.$$

This is precisely $\lambda(W)\lambda(W')$.

There is another way to evaluate the limit. Since the integral is absolutely converge, by Fubini theorem we could rewrite the integral I_c as

$$I_c(W, W') = \int_{\Xi} \int_{N^c \Xi} \Phi_{W_n, W'_n}(x^{-1}g) W_{n+1}(g) \overline{W'_{n+1}}(x) dg dx.$$

Now we could make a substitution $g \mapsto xg$:

$$I_c(W, W') = \int_{\Xi} \int_{\Xi N^c \Xi} \Phi_{W_n, W'_n}(g) W_{n+1}(xg) \overline{W'_{n+1}}(x) dg dx.$$

Again by Fubini theorem we have

$$\begin{aligned} I_c(W, W') &= \int_{\Xi N^c \Xi} \Phi_{W_n, W'_n}(g) \left(\int_{\Xi} W_{n+1}(xg) \overline{W'_{n+1}}(x) dx \right) dg \\ &= \int_{\Xi N^c \Xi} \Phi_{W_n, W'_n}(g) \Phi_{W_{n+1}, W'_{n+1}}(g) dg. \end{aligned}$$

Now by the absolute convergence of

$$\int_{H_n(F)} \Phi_{W_n, W'_n}(g) \Phi_{W_{n+1}, W'_{n+1}}(g) dg,$$

and $\text{supp}(\Phi_{W_{n+1}, W'_{n+1}}) \subset \Xi N_n(F) \Xi$, we conclude that $\lim_{c \rightarrow \infty} I_c(W, W')$ is equal to $\alpha(W, W')$. Finally, we need to verify that under the restriction of support of W, W' , we may still find some choice such that $\lambda(W)\lambda(W')$ does not vanish. But this is clear since we are free to choose Ξ , W_n and W'_n arbitrarily. $\square$

5 The totally definite case

We now prove the part (2) of Theorem 1.2. We hence assume that $G(F_\infty)$ is compact. Equivalently, F is a totally real field, E is a CM extension and the Hermitian spaces W, V are positive definite at every archimedean place v of F . As in the proof of the part (1) of Theorem 1.2, we may assume that the linear functional $\mathcal{P}$ does not vanish on π and $\alpha_v \neq 0$ for all v .

Since $G(F_v)$ is compact for every $v|\infty$, we may choose f_v a function supported in the (relative to H_v) regular semisimple locus. Then there exists a function $f'_v \in G'(F_v)$ which matches f_v (by our convention, this means that for all other isomorphism classes (W_v, V_v) , the functions are chosen to be zero). For non-archimedean places we have no restriction of choice of test functions. For such matching functions f' and f , by Theorem 4.2 we have

$$2^{-2} L(1, \eta)^{-2} I_{\pi_E}(f') = J_\pi(f).$$

For $v|\infty$, as the representation π_v is finite dimensional and $\text{Hom}_{H_v}(\pi_v, \mathbb{C}) \neq 0$, clearly we may choose an f_v supported in the regular semisimple locus so that $J_{\pi_v}(f_v) \neq 0$. Now we may choose $f^\infty = \otimes_{v < \infty} f_v$ so that $J_\pi(f)$ does not vanish. Then for each $v|\infty$, there is a constant $b_{\pi_v} \neq 0$ depending only on π_v such that

$$I_{\Pi_v}(f'_v) = b_{\pi_v} J_{\pi_v}(f_v).$$

Now we don't know how to evaluate b_{π_v} . Nevertheless the same argument as the proof of Prop. 4.4 shows that

$$\frac{|\mathcal{P}(\phi)|^2}{\langle \phi, \phi \rangle_{Pet}} = c_{\pi_\infty} 2^{-2} \mathcal{L} \left(\frac{1}{2}, \pi \right) \prod_v \frac{\alpha_v'^{\natural}(\phi_v, \phi_v)}{\langle \phi_v, \phi_v \rangle_v},$$

where the constant $c_{\pi_\infty} = \prod_{v|\infty} c_{\pi_v}$ and

$$c_{\pi_v} = b_{\pi_v} \kappa_v^{-1} L(1, \eta_v),$$

where κ_v is the constant in Conjecture 4.3.

6 Harmonic analysis on Lie algebra

We establish some basic results to prove the identity between local characters, Theorem 4.5 for a non-split non-archimedean place v .

Relative regular nilpotent elements on $M_{n+1}(F)$. The group H_n , viewed as a subgroup of H_{n+1} , acts on M_{n+1} by conjugation:

$$X^h := hXh^{-1}, \quad X \in M_{n+1}, h \in H_n.$$

Write

$$X = \begin{pmatrix} A & u \\ v & w \end{pmatrix} \in M_{n+1}.$$

The ring of invariants is generated by

$$\mathrm{tr} \wedge^i X, \quad e_{n+1}^* X^j e_{n+1}, \quad 1 \leq i \leq n+1, 1 \leq j \leq n;$$

or

$$\mathrm{tr} \wedge^i A, \quad v A^j u, \quad w, \quad 1 \leq i \leq n, 0 \leq j \leq n-1.$$

The H_n -nilpotent cone $\mathcal{N}$ is defined to be the zeros of all of the above invariant functions on M_{n+1} . An element in M_{n+1} is called H_n -regular (or regular if no confusion arises) if its stabilizer is trivial. Denote

$$\xi_+ = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & \dots & 0 & 1 \\ 0 & \dots & 0 & 0 \end{pmatrix} \in M_{n+1}(F),$$

and ξ_- its transpose. Clearly $\xi_\pm$ are regular nilpotent.

Lemma 6.1. *Let $\xi \in \mathcal{N}$. The following are equivalent.*

1. ξ is regular nilpotent.
2. ξ is H_n -equivalent to ξ_+ .
3. $\Delta_+(\xi) \neq 0$.

Similarly we may replace $+$ by $-$.

Proof. If $X = \begin{pmatrix} A & u \\ v & 0 \end{pmatrix}$ is H_n -nilpotent, we have $A^n = 0$ and $vA^i u = 0$, for $i = 0, 1, \dots, n-1$.

Let r be the dimension of subspace of the column vectors spanned by $A^i u, i = 0, 1, \dots, n-1$, and similarly r' the dimension of the subspace spanned by $vA^i, i = 0, 1, \dots, n-1$. Clearly, we have $r + r' = n$. we need to show that if X is regular unipotent, then either r or r' is equal to n . Indeed, for example if $r = n$, then the column vectors $A^i u, i = 0, 1, \dots, n-1$ form a basis of the n -dimension space. In term of this new basis we see that X becomes ξ_+ .

Suppose $r, r' < n$. Clearly if $r = r' = 0$, X must have positive dimensional stabilizer hence non-regular. Without loss of generality, we may assume that $0 < r < n$. Let L be the subspace spanned by $A^i u, i = 0, 1, \dots, r-1$ (it is easy to see that this is the same as the space spanned by $A^i u, i = 0, 1, \dots, n-1$). And we write the column vector spaces $F^n = L \oplus L'$ for a subspace L' . Then in term of the basis of L given by $A^i u, i = 0, 1, \dots, r-1$, we may write u as $(0, 0, \dots, 1, 0, 0, \dots, 0)^t$ where only the r -th entry is one, and

$$A = \begin{pmatrix} Y & B \\ 0 & Z \end{pmatrix}, \quad Y = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & \dots & 0 & 1 \\ 0 & \dots & 0 & 0 \end{pmatrix} \in M_r(F)$$

Then $A^i u = (0, 0, \dots, 1, 0, 0, \dots, 0)^t$ where only the $(r-i)$ -th entry is one, $i = 0, 1, \dots, r-1$. Hence the conditions $vA^i u = 0$ implies that v is of the form $(0, 0, \dots, 0, *, \dots, *)$ where the first r -entries are all zero.

Now we consider

$$h = \begin{pmatrix} 1_r & \lambda \\ 0 & 1_{n-r} \end{pmatrix} \in \text{GL}_n(F).$$

Clearly the action of h takes X to an element of the same form with B replaced by $B + \lambda Y - Z\lambda$. Hence the stabilizer of X at least contains all h with λ satisfying $\lambda Y - Z\lambda = 0$. Clearly there exists a positive dimensional of such λ since $0 < r < n$. We are done. $\square$

Lemma 6.2. *Let F be any field. Denote by $\mathcal{V}$ the subvariety of $M_{n,F}$ consisting of X that is nilpotent (i.e., $X^n = 0$) of the following form:*

$$X = \begin{bmatrix} * & 1 & 0 & 0 \\ * & * & 1 & 0 \\ \dots & \dots & * & 1 \\ * & \dots & * & * \end{bmatrix} \in M_{n,F}$$

Denote by $\mathcal{V}'$ the subvariety of $M_{n,F}$ consisting of X the same form but with the last row identically zero (not assuming unipotency of X). Then we have a natural projection $p : \mathcal{V} \rightarrow$

$\mathcal{V}'$. Define an F -morphism

$$\begin{aligned}\nu : N_- &\rightarrow \mathcal{V} \\ u &\mapsto u\xi_+u^{-1}\end{aligned}$$

and denote $\nu' = p \circ \nu : N_- \rightarrow \mathcal{V}'$. Then p, ν, ν' are all isomorphisms between affine varieties. Moreover, the Jacobian of ν' is ± 1 .

Here the Jacobian is in term of the obvious coordinates on N_- and $\mathcal{V}'$. Similarly in the proof below. It follows from the lemma that X is conjugate (under a unique element in $N_-(F)$) to ξ_+ . In particular X is regular unipotent (the regularity also follows from the fact that A^ie spans the entire space of column vectors).

Proof. It is easy to see that $u\xi_+u^{-1}$ lies in $\mathcal{V}'$ (this also follows from the proof below). We prove this by induction on n . It is trivial when $n = 1$. We use $N_{n,-}, \mathcal{V}_n$ etc. to indicate the dependence on n . We naturally consider $N_{n-1,-}$ as a sub variety of $N_{n,-}$. We may write $u_n \in N_-$ as a product $u \cdot u_{n-1}$ where $u_{n-1} \in N_{n-1,-}$ and u is of the following form

$$(6.1) \quad u = \begin{bmatrix} & & & 0 \\ & 1_{n-1} & & 0 \\ & & & 0 \\ x_1 & \dots & x_{n-1} & 1 \end{bmatrix}, x_i \in F.$$

Clearly $u_{n-1} = u \cdot u_{n-1}$ defines an isomorphism of affine varieties $N_{n-} = N_{n-1,-} \times F^{n-1}$ with Jacobian ± 1 (in terms of the obvious coordinates).

We write $X_{n-1} = u_{n-1}\xi_{n-1,+}u_{n-1}^{-1} \in \mathcal{V}_{n-1}$, $x = (x_1, \dots, x_{n-1}) \in F^{n-1}$. Then

$$u_n\xi_{n+}u_n^{-1} = u \begin{pmatrix} X_{n-1} & e_{n-1} \\ 0 & 0 \end{pmatrix} u^{-1}$$

is given by

$$\begin{pmatrix} X_{n-1} - e_{n-1}x & e_{n-1} \\ xX_{n-1} - xe_{n-1}x & xe_{n-1} \end{pmatrix}.$$

Replacing x by $-x + x_{n-1}$ where x_{n-1} is the last row of X_{n-1} , by induction hypothesis that ν'_{n-1} is an isomorphism, we conclude that so is ν'_n . The formula also shows the Jacobian of ν'_n is ± 1 by induction hypothesis.

It remains to show that p_n is an isomorphism. We will define an inverse of p_n inductively. We start with X of the form

$$\begin{bmatrix} & & & 0 \\ & X_{n-1} & & 0 \\ & & & 1 \\ a_1 & \dots & a_{n-1} & a_n \end{bmatrix}, X_{n-1} \in \mathcal{V}_{n-1}.$$

We consider a unipotent u in $N_-(F)$ of the form as in (6.1). Then we see that the most lower right entry of uXu^{-1} is given by $a_n + x_{n-1}$. Thus we may choose $x_{n-1} = -a_n$ and hence assume that in X we have $a_n = 0$ and in u we have $x_{n-1} = 0$. Then inductively if we have $a_n = a_{n-1} = \dots = a_{n-i} = 0$, and $x_{n-1} = x_{n-2} = \dots = x_{n-i-1} = 0$, then (using the property of X_{n-1}) we see that the $(n, n-i-1)$ -th entry of uXu^{-1} is given by $a_{n-i-1} + x_{n-i-2}$. Therefore we have (a unique such) u of the above form such that uXu^{-1} is of the form

$$uXu^{-1} = \begin{bmatrix} & & & 0 \\ & X'_{n-1} & & 0 \\ & & & 1 \\ a'_1 & 0 & \dots & 0 \end{bmatrix}, \quad a'_1 \in F,$$

where X'_{n-1} retains the same shape as X_{n-1} (by computation). Note that

$$\det(X) = \pm(a_1 - a_n * \det(X_{n-1})).$$

Now it is easy to see that $0 = \det(X) = \det(uXu^{-1}) = \pm(a'_1 - 0)$. Hence $a'_1 = 0$. Moreover the unipotence of gXg^{-1} (being block upper triangular) now implies the unipotence of X'_{n-1} . Now it is clearly that by induction hypothesis the above process defines an inverse of p_n . $\square$

We also have an similar but easier statement on the upper unipotent N_+ .

Lemma 6.3. *Let F be any field. Denote by $\mathcal{V}_+$ the subvariety of $M_{n,F}$ consisting of X that is nilpotent (i.e., $X^n = 0$) of the following form:*

$$X = \begin{bmatrix} 0 & 1 & * & * \\ 0 & 0 & 1 & * \\ \dots & \dots & 0 & 1 \\ 0 & \dots & 0 & 0 \end{bmatrix} \in M_{n,F}$$

Define an F -morphism

$$\begin{aligned} \nu_+ : N_+ &\rightarrow \mathcal{V}_+ \\ u &\mapsto u\xi_+u^{-1}. \end{aligned}$$

Then ν_+ is an isomorphisms between affine varieties. Moreover, the Jacobian of ν_+ is ± 1 .

Proof. Similar to the previous one. We omit the detail. $\square$

Regular nilpotent orbital integral. Let F be a local field. We now define the (H_n, η) -orbital integral of a regular nilpotent orbit. For $X = \begin{pmatrix} A & u \\ v & w \end{pmatrix} \in M_{n+1}(F)$, we define a matrix

$$\delta_+(X) := (A^{n-1}u, A^{n-2}u, \dots, u) \in M_n(F),$$

and its determinant

$$\Delta_+(X) = \det(\delta_+(X)).$$

Similarly, we define

$$\delta_-(X) := \delta_+(X^t), \quad \Delta_-(X) = \det(\delta_-(X)),$$

and

$$\Delta := \Delta_+ \Delta_-.$$

Since the orbits of $\xi_\pm$ are not closed, we need to regularize the orbital integral. We consider for $s \in \mathbb{C}$:

$$O(X, f, s) = \eta(\Delta_+(X)) \int_{H_n(F)} f(X^h) \eta(h) |\det(h)|^s dh.$$

Lemma 6.4. *The integral $O(\xi_\pm, f, s)$ converges absolutely when $\operatorname{Re}(s) > 1 - \frac{1}{n}$ and extends to a meromorphic function on $s \in \mathbb{C}$ with at most a simple pole at $s = 1 - 1/i$, for even integers i with $1 < i \leq n$.*

Proof. We use the Iwasawa decomposition of $H_n(F) = KAN$. Define

$$f_K(X) = \int_K f(kXk^{-1}) dk.$$

By Iwasawa decomposition on $H_n(F)$, we have

$$\int_{A(F)} \int_{N(F)} f_K(au\xi_+u^{-1}a^{-1}) \eta(a) |a|^s |\delta(a)| du da,$$

where the modular character

$$\delta(a) = a_1^{n-1} a_2^{n-3} \dots a_n^{-(n-1)}, \quad a = \operatorname{diag}[a_1, \dots, a_n].$$

By Lemma 6.3, this is

$$\int_{A(F)} \int_{\mathcal{V}_+(F)} f_K(axa^{-1}) \eta(a) |a|^s dx da$$

Replacing x_{ij} by $x_{ij} a_j a_i^{-1}$, $1 \leq i \leq j-2 \leq (n+1)-2$ and setting $a_{n+1} = 1$, we may cancel partially the factor $|\delta(a)|$:

$$\int_{A(F)} \int_{\mathcal{V}_+(F)} f_K \begin{bmatrix} 0 & \frac{a_1}{a_2} & x_{13} & x_{14} & * \\ 0 & 0 & \frac{a_2}{a_3} & * & * \\ 0 & 0 & 0 & \dots & \dots \\ \dots & \dots & 0 & 0 & a_n \\ 0 & \dots & 0 & 0 & 0 \end{bmatrix} \eta(a) |a|^s |a_2 a_3 \dots a_n|^{-1} dx da$$

Substitute $b_i := a_i/a_{i+1}, i = 1, 2, \dots, n-1, b_n = a_n$:

$$\int_{A(F)} \int_{\mathcal{V}_+(F)} f_K \begin{bmatrix} 0 & b_1 & x_{13} & x_{14} & * \\ 0 & 0 & b_2 & * & * \\ 0 & 0 & 0 & \dots & \dots \\ \dots & \dots & 0 & 0 & b_n \\ 0 & \dots & 0 & 0 & 0 \end{bmatrix} \eta(b_1 b_3 \dots) \left| \prod_{i=1}^n b_i^i \right|^{-1+s} \prod_{i=1}^n db_i dx.$$

Now it is clear that the integral converges absolutely if $\operatorname{Re}(i(-1+s)) > -1$ for all $i = 1, 2, \dots, n$, or equivalently $\operatorname{Re}(s) > 1 - \frac{1}{n}$. And by Tate's local zeta integral, the integral extends meromorphically to $s \in \mathbb{C}$ with at most simple poles at those s such that

$$i(-1+s) = -1, \quad i = 2, 4, \dots, 2[n/2].$$

Namely $s = 1 - 1/i$, for even i with $1 < i \leq n$.

□

Definition 6.5. For $f \in \mathcal{C}_c^\infty(M_{n+1})$, we define the regular nilpotent orbital integral $O(\xi_\pm, f)$, also denoted by $\mu_{\xi_\pm}(f)$ as

$$\mu_{\xi_\pm}(f) = O(\xi_\pm, f) := O(\xi_\pm, f, 0).$$

This defines an (H_n, η) -invariant distribution on M_{n+1} .

A regular section of π . Denote

$$M_{n+1,+} := \{X \in M_{n+1} \mid \Delta_+(X) \neq 0\}$$

Then $X \in M_{n+1,+}$ if and only if $\delta_+(X) \in \operatorname{GL}_{n,F}$, or equivalently $u, Au, \dots, A^{n-1}u$ are linearly independent. The latter property implies that the stabilizer (under the H_n -action) of any $X \in M_{n+1,+}$ must be trivial. Indeed, if $hXh^{-1} = X$, we have $hAh^{-1} = A, hu = u$. Therefore $hA^i u = hA^i h^{-1} \cdot hu = A^i u$ for all $i = 0, 1, \dots, n-1$. It follows that $h = 1_n$ from the linear independence of $u, Au, \dots, A^{n-1}u$.

We will write $\mathcal{X} = \mathbb{A}^n \times \mathbb{A}^{n+1}$,⁶ the affine space of dimension $2n+1$. Then the second set of generators defines a morphism that is constant on H_n -orbits

$$\begin{aligned} \pi : M_{n+1} &\rightarrow \mathcal{X} = \mathbb{A}^n \times \mathbb{A}^{n+1}, \\ \begin{pmatrix} A & u \\ v & w \end{pmatrix} &\mapsto (a, b), \end{aligned}$$

⁶we only use here $\mathbb{A}$ to denote the affine line.

where $a = (a_1, \dots, a_n)$, $b = (b_0, \dots, b_n)$, $a_{i+1} = \text{tr } \wedge^{i+1} A$, $b_0 = w$ and $b_i = vA^i u$ for $0 \leq i \leq n-1$. Now we define a section of the morphism $\pi : M_{n+1} \rightarrow \mathcal{X}$:

$$\sigma : \mathcal{X} \rightarrow M_{n+1}$$

$$(a, b) \mapsto \begin{pmatrix} a_1 & 1 & 0 & 0 & 0 \\ a_2 & 0 & 1 & 0 & 0 \\ \dots & 0 & 0 & 1 & 0 \\ a_n & 0 & 0 & 0 & 1 \\ b_n & \dots & \dots & b_1 & b_0 \end{pmatrix}.$$

We note that ξ_+ is precisely the image of $0 \in \mathcal{X}$ under σ .

Lemma 6.6. *The morphism σ is a section of π , i.e.:*

$$\sigma \circ \pi = \text{id}.$$

The image of σ lies in $M_{n+1,+}$ (in particular, σ is a regular section, in the sense that the image $\sigma(a, b)$ is always H_n -regular).

Proof. Indeed it is easy to check that

$$\det \left(T + \begin{pmatrix} a_1 & 1 & 0 & 0 \\ a_2 & 0 & 1 & 0 \\ \dots & 0 & 0 & 1 \\ a_n & 0 & 0 & 0 \end{pmatrix} \right) = T^n + a_1 T^{n-1} - a_2 T^{n-2} + \dots + (-1)^{n-1} a_n,$$

and the b -invariants of $\sigma(a, b)$ are $(b_0, b_1, b_2, \dots, b_n)$. This shows that σ is a section of π . To see that the image of σ lies in $M_{n+1,+}$, we note that for any $(a, b) \in \mathcal{X}$ we have

$$\delta_+(\sigma(a, b)) = 1_n.$$

□

Note that δ_+ (restricted to $M_{n+1,+}$) defines a morphism

$$\delta_+ : M_{n+1,+} \rightarrow \text{GL}_n$$

$$X \mapsto \delta_+(X).$$

Moreover, it has the property

$$\delta_+(hXh^{-1}) = h\delta_+(X), \quad h \in \text{GL}_{n,F}.$$

Proposition 6.7. *We define an H_n -equivariant morphism*

$$\iota : \text{GL}_n \times \mathcal{X} \rightarrow M_{n+1,+}$$

$$(h, (a, b)) \mapsto h\sigma(a, b)h^{-1},$$

where H_n -acts on LHS by left translation on the first factor, trivially on the second. Then ι is an isomorphism with its inverse given by $(\delta_+, \pi|_{\mathfrak{s}_+})$.

Proof. It suffices to prove that $(\delta_+, \pi) \circ \iota = id$ and $\iota \circ (\delta_+, \pi) = id$. To show the first identity we note that the invariants of $h\sigma(a, b)h^{-1}$ (being the same as $\sigma(a, b)$) are (a, b) . Hence it is enough to show that $\delta_+(\iota(h, (a, b))) = h$. This follows from the fact that $\delta_+(\sigma(a, b)) = 1_n$ and $\delta_+(hXh^{-1}) = h\delta_+(X)$. Now we show the second identity. Let $X = \begin{pmatrix} A & u \\ v & w \end{pmatrix} \in M_{n+1,+}$. Let $(a, b) = \pi(X)$ be the invariants of X and $h = \delta_+(X) = (A^{n-1}u, A^{n-2}u, \dots, u)$. Denote $\iota \circ (\delta_+, \pi)(X) = \begin{pmatrix} A' & u' \\ v' & w' \end{pmatrix}$. Clearly $w = w'$. By the first identity, the elements $\iota \circ (\delta_+, \pi)(X)$ and X have the same invariants. In particular,

$$\det(T + A) = T^n + \sum_{i=1}^n (-1)^{i-1} a_i T^{n-i},$$

and

$$A^n = \sum_{i=1}^n a_i A^{n-i}.$$

This implies that

$$A\delta_+(X) = (A^n u, A^{n-1}u, \dots, Au) = \delta_+(X) \begin{pmatrix} a_1 & 1 & 0 & 0 \\ a_2 & 0 & 1 & 0 \\ \dots & 0 & 0 & 1 \\ a_n & 0 & 0 & 0 \end{pmatrix}.$$

Since $\delta_+(X) = h$ is invertible, we obtain

$$A = h \begin{pmatrix} a_1 & 1 & 0 & 0 \\ a_2 & 0 & 1 & 0 \\ \dots & 0 & 0 & 1 \\ a_n & 0 & 0 & 0 \end{pmatrix} h^{-1} = A'.$$

Obviously we have $u = \delta_+(X)e_n = he_n = u'$ ($e_n = (0, 0, \dots, 0, 1)^t$). Finally since $b_i = vA^{i-1}u$, $(b_n, b_{n-1}, \dots, b_1) = v\delta_+(X)$, we have

$$v = (b_n, b_{n-1}, \dots, b_1)\delta_+(X)^{-1} = (b_n, b_{n-1}, \dots, b_1)h^{-1} = v'.$$

This completes the proof of the second identity. $\square$

Now we apply the results above to the orbital integrals. Assume that F is a non-archimedean local field.

Corollary 6.8. *The intersection $M_{n+1,+}(F) \cap \mathcal{N}$ is equal to the H_n -orbit of ξ_+ . In particular, for a function $f \in \mathcal{C}_c^\infty(M_{n+1}(F))$ supported on $M_{n+1,+}(F)$, any distribution on $M_{n+1}(F)$ supported in the closed subset $\mathcal{N} \setminus (H_n \cdot \xi_+)$ of $M_{n+1}(F)$ vanishes on f .*

Proof. Since ξ_+ is precisely the image of $0 \in \mathcal{X}$ under σ , the H_n -orbit of ξ_+ is then the image of $H_n \times \{0\}$ under ι . We also have $M_{n+1,+} \cap \mathcal{N} = (\pi|_{M_{n+1,+}})^{-1}(0)$, the fiber of $0 \in \mathcal{X}$ under $\pi|_{M_{n+1,+}}$. By Proposition above the fiber $(\pi|_{M_{n+1,+}})^{-1}(0)$ is precisely the image of $H_n \times \{0\}$ under ι . This proves the first assertion. The ‘‘In particular’’ part is clearly from the definition of support of a distribution. $\square$

Proposition 6.9. *For any $f \in \mathcal{C}_c^\infty(M_{n+1,+}(F)) \subset \mathcal{C}_c^\infty(M_{n+1}(F))$, the orbital integral*

$$\phi_f(x) := O(\sigma(x), f)$$

defined for regular semisimple $x \in \mathcal{X}$ extends to a locally constant function with compact support on $\mathcal{X}$ (i.e., $\phi_f \in \mathcal{C}_c^\infty(\mathcal{X})$). And conversely, given any function ϕ in $\mathcal{C}_c^\infty(\mathcal{X})$, there exists $f \in \mathcal{C}_c^\infty(M_{n+1,+}(F))$ such that $O(\sigma(x), f) = \phi(x)$ for all regular semisimple x .

Proof. Since $\Delta_+(\sigma(x)) = 1$, the normalized orbital integral is given by

$$O(\sigma(x), f) = \int_{H_n} f(h\sigma(x)h^{-1})\eta(h)dh.$$

By the H_n -equivariant isomorphism $\iota : H_n(F) \times \mathcal{X}(F) \rightarrow M_{n+1,+}(F)$, corresponding to f we have an element denoted by f' in $\mathcal{C}_c^\infty(H_n \times \mathcal{X})$ with the property that

$$O(\sigma(x), f) = \int_{H_n} f'(h, x)\eta(h)dh.$$

The integral in RHS is clearly convergent absolutely for all $x \in \mathcal{X}$ and defines an element in $\mathcal{C}_c^\infty(\mathcal{X})$. The converse is clearly now by the isomorphism ι . $\square$

Remark 7. The proof also shows that if f is supported in $M_{n+1}(F)$, the integral $O(\xi_+, f, s)$ converges absolutely for any $s \in \mathbb{C}$.

Harmonic analysis on $\mathfrak{s}_{n+1}$. We also need to consider the induced H_n -action on the tangent space $\mathfrak{s}_{n+1}$ at 1 of the symmetric space S_{n+1} :

$$\mathfrak{s}_{n+1}(F) = \{X \in M_{n+1}(E) | X + \overline{X} = 0\}.$$

For any choice of non-zero $\tau \in E^-$, we have an isomorphism

$$M_{n+1}(F) \simeq \mathfrak{s}_{n+1}(F),$$

defined by $X \mapsto \tau X$. In particular, we will abuse the notation $\xi_\pm$ to denote $\tau\xi_\pm$ if we want to consider the regular unipotent orbit on $\mathfrak{s}_{n+1}$. Then clearly every result established in this section holds for $\mathfrak{s}_{n+1}$.

7 Smoothing local periods

Convolutions. We introduce some abstract notions for the use of this and the next section. Let F be a p -adic local field and G the F -points of some reductive group. We consider the space of $\mathcal{C}_c^\infty(G)$ with an (anti-) involution $*$ defined by:

$$f^*(g) := \overline{f(g^{-1})}, \quad f \in \mathcal{C}_c^\infty(G).$$

We will also use the other (anti-) involution defined by

$$f^\vee(g) = f(g^{-1}).$$

Let dg be a Haar measure on G . Let H be a closed subgroup of G , *unimodular* with a Haar measure dh . We define a left and a right action of $\mathcal{C}_c^\infty(H)$ on $\mathcal{C}_c^\infty(G)$ as follows: for $f \in \mathcal{C}_c^\infty(G)$ and $\phi \in \mathcal{C}_c^\infty(H)$, we define *convolutions* $f \circ \phi$ and $\phi \circ f$ both in $\mathcal{C}_c^\infty(G)$:

$$(f \circ \phi)(g) = \int_H f(gh^{-1})\phi(h)dh$$

and

$$(\phi \circ f)(g) = \int_H \phi(h)f(h^{-1}g)dh = \int_H \phi(h^{-1})f(hg)dh.$$

This also applies to the case $H = G$. Then we have

$$(f \circ \phi)^* = \phi^* \circ f^*, \quad (\phi \circ f)^* = f^* \circ \phi^*.$$

If we have two closed unimodular subgroups H_1, H_2 , we could iterate the definition: for example, for $f \in \mathcal{C}_c^\infty(G)$ and $\phi_i \in \mathcal{C}_c^\infty(H_i)$, we define

$$\phi_1 \circ \phi_2 \circ f := \phi_1 \circ (\phi_2 \circ f) \in \mathcal{C}_c^\infty(G).$$

Now if we have a smooth representation π of G (hence its restriction to H is a smooth representation as well), as usual we define $\pi(f)$ and $\pi(\phi)$ to be the endomorphisms of π :

$$\pi(f) = \int_G f(g)\pi(g)dg, \quad \pi(\phi) = \int_H \phi(h)\pi(h)dh.$$

Then we have $\pi(f \circ \phi) = \pi(f)\pi(\phi) \in \text{End}(\pi)$ and so on. If π has an G -invariant inner product $\langle \cdot, \cdot \rangle$, we then have

$$\langle \pi(f)u, u' \rangle = \langle u, \pi(f^*)u' \rangle, \quad u, u' \in \pi.$$

The question addressed in this section has the following shape. Let π be a smooth admissible representation of G with an invariant inner product. Very often the algebraic dual space $\pi^* := \text{Hom}(\pi, \mathbb{C})$ is much larger than the congragredient $\tilde{\pi}$ (the subspace of π^* consisting of smooth linear functional, i.e., K -finite vectors in π^* for some open compact K). Very often we will be interested in some distinguished element called ℓ in $\pi^* \setminus \tilde{\pi}$ (the set of non-smooth

linear functionals). It is also very often useful to find some $\phi \in \mathcal{C}_c^\infty(H)$, for suitable subgroup H (smaller than G in order to be useful), such that $\pi^*(\phi^*)\ell$ is indeed smooth (and non-zero), i.e., there exists $0 \neq u' \in \pi$ such that for all $u \in \pi$:

$$\ell(\pi(\phi)u) = \langle u, u' \rangle.$$

We will encounter this kind of questions for the local Flicker-Rallis period, and the local Rankin-Selberg period.

A compactness lemma. Now we return to our setting. Let E/F be a quadratic extension of non-archimedean local fields of characteristic zero. Let η be the quadratic character associated to E/F , and denote $\eta_n = \eta^{n-1}$. Let $\varphi_{n-1} \in \mathcal{C}_c^\infty(H_{n-1}(E))$ and $\phi_{n-1} \in \mathcal{C}_c^\infty(M_{n-1,1}(E))$. We consider the Fourier transform of ϕ_{n-1} as a function on the $M_{1,n-1}(E)$ by

$$\widehat{\phi}_{n-1}(X) = \int_{M_{1,n-1}(E)} \phi_{n-1}(Y) \psi_E(\text{tr}(XY)) dY.$$

To the pair $(\varphi_{n-1}, \phi_{n-1})$ we associate a new function on $H_{n-1}(E)$ by

$$(7.1) \quad \widetilde{W}_{\varphi_{n-1}, \phi_{n-1}}(g) := \widehat{\phi}_{n-1}(-e_{n-1}^* g) \int \varphi_{n-1}(g^{-1} u \epsilon_{n-1} h) \overline{\psi}_E(u) \eta_n(h) du dh,$$

where $u \in N_{n-1}(E)$, $h \in N_{n-1}(F) \backslash H_{n-1}(F)$. Clearly we have

$$W_{\varphi_{n-1}, \phi_{n-1}}(ug) = \overline{\psi}_E(u) W_{\varphi_{n-1}, \phi_{n-1}}(g)$$

for $u \in N_{n-1}(E)$.

We would like to obtain a function with support compact modulo $N_{n-1}(E)$ by imposing some conditions on $\varphi_{n-1}, \phi_{n-1}$. We construct a subspace of special functions by the following procedure. Fix integers $k \geq 0, m > 0$. Define a sublattice of $\mathcal{O}_E$

$$(7.2) \quad \Lambda = \Lambda_k := \varpi^k \mathcal{O}_E.$$

It decomposes as $\Lambda_k = \Lambda_k^+ \oplus \Lambda_k^-$ where $\Lambda_k^\pm = \varpi^k \mathcal{O}_{E^\pm}$. Then the Fourier transform of $1_\Lambda \in \mathcal{C}_c^\infty(E)$ is a non-zero multiple of 1_{Λ^*} for a lattice $\Lambda^* \subset E$ (depending on ψ). We naturally view $\mathbb{C}[\varpi^m \Lambda / \varpi^{2m} \Lambda]$ as a subspace of $\mathcal{C}_c^\infty(E)$ consisting of functions supported in $\varpi^m \Lambda$ and invariant by $\varpi^{2m} \Lambda$. We define a “dagger” space of level (k, m) , denote by $\mathbb{C}[\varpi^m \Lambda_k / \varpi^{2m} \Lambda_k]^\dagger$ or $\mathcal{C}_c^\infty(E)_{k,m}^\dagger$, as a subspace of $\mathbb{C}[\varpi^m \Lambda / \varpi^{2m} \Lambda]$ spanned by functions $\theta = \theta^+ \otimes \theta^-, \theta^\pm \in \mathcal{C}_c^\infty(E^\pm)$ satisfying:

- θ^+ is a multiple of $1_{\varpi^m \Lambda_k^+}$.
- The Fourier transform $\widehat{\theta} \in \mathbb{C}[\varpi^{-2m} \Lambda^* / \varpi^{-m} \Lambda^*]$ is (non-emptyly) supported in $\varpi^{-2m} \Lambda^* - \varpi^{-2m+1} \Lambda^*$. Equivalently, $\widehat{\theta^-}$ is supported in $\varpi^{-2m} \Lambda^{*-} - \varpi^{-2m+1} \Lambda^{*-}$ where $\Lambda^{*-} = \Lambda^* \cap E^-$.

In particular, every element in $\mathbb{C}[\varpi^m \Lambda / \varpi^{2m} \Lambda]^\dagger$ is invariant under multiplication by $1 + \varpi^m \mathcal{O}_E$. We also denote by $\mathcal{C}_c^\infty(M_{n-1,1}(E))_{k,m}^\dagger$ the space of functions on $M_{n-1,1}(E)$ of the form $\phi_{n-1} = \bigotimes_{1 \leq i \leq n} \phi^i$ in the way that $\phi_{n-1}(x) = \prod_i \phi^i(x_i)$ if $x = (x_1, \dots, x_{n-1})^t \in M_{n-1,1}(E)$ satisfying

- When $1 \leq i \leq n-1$, ϕ^i is the characteristic function of $\varpi^m \Lambda$; ϕ^n is an element of $\mathbb{C}[\varpi^m \Lambda / \varpi^{2m} \Lambda]^\dagger$.

We denote by $\mathcal{C}_c^\infty(H_{n-1}(E))_{k,m}^\dagger$ the space of functions on $H_{n-1}(E)$ of the form $\varphi_{n-1} = \bigotimes_{1 \leq i,j \leq n-1} \varphi^{ij}$ in the way that $\varphi_{n-1}(g) = \prod_{i,j} \varphi^{ij}(g_{ij})$ if $g = (g_{ij})$ satisfying:

- When $1 \leq i, j \leq n-1$, $j-i \neq 0, 1$, φ^{ij} is the characteristic function of $\varpi^m \Lambda$.
- When $1 \leq i = j \leq n-1$, φ^{ij} is the characteristic function of $1 + \varpi^m \Lambda$.
- When $1 \leq i = j-1 \leq n-2$, φ^{ij} is an element of $\mathbb{C}[\varpi^m \Lambda / \varpi^{2m} \Lambda]^\dagger$.

Let $\mathcal{C}_c^\infty(M_{n-1,1}(E))_m^\dagger$ be the union of $\mathcal{C}_c^\infty(M_{n-1,1}(E))_{k,m}^\dagger$ for all $k \geq 0$. Similarly for $\varphi_{n-1} \in \mathcal{C}_c^\infty(H_{n-1}(E))_m^\dagger$. We say that the pair $(\varphi_{n-1}, \phi_{n-1})$ is *m-admissible* if $\phi_{n-1} \in \mathcal{C}_c^\infty(M_{n-1,1}(E))_m^\dagger$, and $\varphi_{n-1} \in \mathcal{C}_c^\infty(H_{n-1}(E))_m^\dagger$. We also call them “dagger” space of level m .

The pair $(\varphi_{n-1}, \phi_{n-1})$ defines a function denoted by $\varphi_{n-1} \otimes \phi_{n-1}$ on the mirabolic group P_n of $H_n(E)$ with support in $1 + \mathfrak{p}_n(\varpi^m \Lambda_k)$ where $\mathfrak{p}_n$ is the Lie algebra of P_n :

$$\varphi_{n-1} \otimes \phi_{n-1} \left[\begin{pmatrix} x & \\ & 1 \end{pmatrix} \begin{pmatrix} 1_{n-1} & u \\ & 1 \end{pmatrix} \right] = \varphi_{n-1}(x) \phi_{n-1}(u).$$

Some property of admissible functions will not be used until next section. Our key lemma of this section is as follows.

Lemma 7.1. *Assume that $(\varphi_{n-1}, \phi_{n-1})$ is m-admissible for some $m > 0$.*

- (1) *Then the support of the function $\widetilde{W}_{\varphi_{n-1}, \phi_{n-1}}$ is compact modulo $N_{n-1}(E)$, i.e., it defines an element in*

$$\mathcal{C}_c^\infty(N_{n-1}(E) \backslash H_{n-1}(E), \overline{\psi}_E).$$

More precisely, $\widetilde{W}_{\varphi_{n-1}, \phi_{n-1}}(\epsilon_{n-1}g)$ is nonzero only when

$$g \in H'_{n-1}(E) = N_{n-1}(E)A_{n-1}(E)N_{n-1,-}(E).$$

- (2) *For $i = n-2, \dots, 1$, we define recursively ϕ_i, ϕ'_{i+1} such that*

$$\varphi_{i+1} = \varphi_i \otimes \phi_i \otimes \phi'_{i+1}$$

where

$$\varphi_i \in \mathcal{C}_c^\infty(M_{i,i}(E)), \phi_i \in \mathcal{C}_c^\infty(M_{i,1}(E)), \phi'_{i+1} \in \mathcal{C}_c^\infty(M_{1,i+1}(E)).$$

Set $\phi'_1 = \varphi_1$ so that we have the following pattern of decomposition of $\varphi_{n-1} \otimes \phi_{n-1}$:

ϕ'_1	ϕ_1	ϕ_2	$\cdots$	ϕ_{n-1}
ϕ'_2				
$\vdots$				
ϕ'_{n-1}				

Set $\phi' = \otimes_{i=1}^{n-1} \phi'_i$ viewed as a function on $B_{n-1,-}(E)$ or $\mathfrak{b}_{n-1,1}(E)$. Denote by $d_n = n(n-1)(n-2)/6$. Then the value of $\widetilde{W}_{\varphi_{n-1}, \phi_{n-1}}(\epsilon_{n-1}g)$ when $g = yz \in A_{n-1}N_{n-1,-}(F)$ for

$$y = \begin{pmatrix} y_1 y_2 \cdots y_{n-1} & & & \\ & \ddots & & \\ & & y_1 y_2 & \\ & & & y_1 \end{pmatrix} \in A_{n-1}(F),$$

and

$$z = \prod_{i=1}^{n-2} \begin{pmatrix} 1_i & \\ z_i & 1 \end{pmatrix} \in N_{n-1,-}(F), \quad z_i \in M_{1,i}(F),$$

is given by the product of a constant

$$|\tau|_E^{d_n} \left(\int_{B_{n-1,-}(F)} \phi'(b) db \right)$$

where $\tau^{d_n} = \delta_{n-1}(\epsilon_{n-1}) = \det(\text{Ad}(\epsilon_{n-1}) : N_{n-1}(E))$, and

$$(7.3) \quad \eta_n(y) |\delta_{n-1}(y)|_F \prod_{i=1}^{n-1} \widehat{\phi_{n-i}}(-y_i(z_{n-i-1}, 1)\tau).$$

Here the measure db on $B_{n-1,-}(F)$ is either the left or the right invariant one; they give the same value to the integral since the support of ϕ' is in $1 + \varpi \mathfrak{b}_{n-1,-}(\mathcal{O}_E)$.

Proof. It suffices to consider the following function:

$$w(g) = \int_{N_{n-1}(F) \backslash H_{n-1}(F)} \int_{N_{n-1}(E)} \varphi(g^{-1}uh) \overline{\psi}_\tau(u) \eta_n(h) du dh,$$

where

$$\psi_\tau(u) = \psi_{E,\tau}(u) = \psi_E(\epsilon_{n-1} u \epsilon_{n-1}^{-1}).$$

Indeed, by a suitable substitution we have

$$\widetilde{W}_{\varphi, \phi_{n-1}}(\epsilon_{n-1}g) = |\tau|_E^{d_n} \widehat{\phi_{n-1}}(-e_{n-1}^* \tau g) w(g).$$

By the condition on $\hat{\phi}_{n-1}$, $\hat{\phi}_{n-1}(e_{n-1}^*g)$ is zero unless the $(n-1, n-1)$ -th entry of g is nonzero. Up to the $N_{n-1}(E)$ -action, such $g \in H_{n-1}(E)$ is of the form:

$$g = y_1 \begin{pmatrix} x_{n-2} & \\ & 1 \end{pmatrix} \begin{pmatrix} 1_{n-2} & \\ z_{n-2} & 1 \end{pmatrix}, y_1 \in E^\times, x_{n-2} \in H_{n-2}(E), z_{n-2} \in M_{1,n-2}(E).$$

By the support condition on $\hat{\phi}_{n-1}$, for $w(g)$ to be nonzero, $|y_1|$ lies in a compact region of $E^\times$ and $\|z_{n-2}\|$ is small enough so that φ is invariant under left multiplication by $\begin{pmatrix} 1_{n-2} & 0 \\ z_{n-2} & 1 \end{pmatrix}$. Hence we have

$$(7.4) \quad \widetilde{W}_{\varphi_{n-1}, \phi_{n-1}}(\epsilon_{n-1}g) = |\tau|_E^{d_n} \hat{\phi}_{n-1}(\tau y_1(z_{n-2}, 1)) w \left[y_1 \begin{pmatrix} x_{n-2} & \\ & 1 \end{pmatrix} \right].$$

Therefore it is enough to consider $w(g)$ when $g = y_1 \begin{pmatrix} x_{n-2} & \\ & 1 \end{pmatrix}$ for $x_{n-2} \in H_{n-2}(E)$.

The integration on $h \in H_{n-1}(F)$ can be replaced by the integration over the open dense subset $H'_{n-1}(F) = N_{n-1}(F)A_{n-1}(F)N_{n-1,-}(F)$. We write $h \in H'_{n-1}(F)$ as

$$h = b_1 \begin{pmatrix} 1_{n-2} & c'_{n-2} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a_{n-2} & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1_{n-2} & 0 \\ c_{n-2} & 1 \end{pmatrix},$$

where $b_1 \in F^\times$, $a_{n-2} \in H'_{n-2}(F)$, $c_{n-2} \in M_{1,n-2}(F)$, $c'_{n-2} \in M_{n-2,1}(F)$. The measure on $H'_{n-1}(F)$ can be decomposed as

$$dh = |a_{n-2}|^{-1} |b_1|^{-1} db_1 dc_{n-2} da_{n-2} dc'_{n-2}.$$

We write

$$u = \begin{pmatrix} 1_{n-2} & v_{n-2} \\ & 1 \end{pmatrix} \begin{pmatrix} u_{n-2} & \\ & 1 \end{pmatrix} \in N_{n-1}(E).$$

In the decomposition of h , we may ignore the factor in $N_{n-1}(F)$. Then the product $g^{-1}uh$ is equal to

$$b_1 y_1^{-1} \begin{pmatrix} x_{n-2}^{-1} & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1_{n-2} & v_{n-2} \\ & 1 \end{pmatrix} \begin{pmatrix} u_{n-2} & \\ & 1 \end{pmatrix} \begin{pmatrix} a_{n-2} & \\ & 1 \end{pmatrix} \begin{pmatrix} 1_{n-2} & \\ c_{n-2} & 1 \end{pmatrix}.$$

Then the last row of the product is equal to $y_1^{-1}b_1(c_{n-2}, 1) \in M_{1,n-1}(E)$. By the condition on ϕ'_{n-1} , we can assume the $\|c_{n-2}\|$ is small enough so that φ_{n-1} is invariant under right translation by $\begin{pmatrix} 1_{n-2} & \\ c_{n-2} & 1 \end{pmatrix}$. The product of the first four matrices is then

$$y_1^{-1}b_1 \begin{pmatrix} x_{n-2}^{-1}u_{n-2}a_{n-2} & x_{n-2}^{-1}v_{n-2} \\ 0 & 1 \end{pmatrix}.$$

The integrations on c_{n-2} and v_{n-2} yield respectively

$$\int_{M_{n-2,1}(F)} \phi'_{n-1}(y_1^{-1}b_1(c_{n-2}, 1)) dc_{n-2},$$

$$\int_{M_{n-2,1}(E)} \phi_{n-2}(b_1 y_1^{-1} x_{n-2}^{-1} v_{n-2}) \overline{\psi}_\tau(v_{n-2}) dv_{n-2} = |b_1^{-1} y_1|_E^{n-2} |x_{n-2}|_E \widehat{\phi}_{n-2}(-e_{n-2} \tau y_1 b_1^{-1} x_{n-2}).$$

(here we note that the Fourier transform of ϕ_{n-2} is defined by the character ψ_E .) Therefore $w(g)$ is equal to the integration of the function of $b_1 \in F^\times$ given by the product of the above two terms and

$$\int_{N_{n-2}(F) \setminus H_{n-2}(F)} \int_{N_{n-2}(E)} \varphi_{n-2}(y_1^{-1} b_1 x_{n-2}^{-1} u_{n-2} a_{n-2}) \overline{\psi}_\tau(u_{n-2}) |a_{n-2}|^{-1} \eta(h) du_{n-2} da_{n-2}$$

with respect to the measure $|b_1|^{(n-1)s} d^* b_1$. We may repeat the process and hence may assume that the function on $H_{n-2}(E)$ mapping g_{n-2} to

$$\widehat{\phi}_{n-2}(-e_{n-2} g_{n-2}) \int_{N_{n-2}(F) \setminus H_{n-2}(F)} \int_{N_{n-2}(E)} \varphi_{n-2}(g_{n-2}^{-1} u_{n-2} a_{n-2}) \overline{\psi}_\tau(u_{n-2}) |a_{n-2}|^{-1} \eta_n(a) du_{n-2} da_{n-2}$$

is nonzero unless $g_2 \in H'_{n-2}(E)$ and its support is compact modulo $N_{n-2}(E)$. As we know that y_1 is in a compact region of $E^\times$, the integration of b_1 must also be in a compact region. This implies that $w(g) \neq 0$ only when $x_{n-2} \in H'_{n-2}(E)$ and in a region compact modulo $N_{n-2}(E)$. By the boundedness of z_{n-2} as shown in the equation (7.4), we complete the proof of the part (1).

To show the part (2), we keep track of the computation above. Since we are now assuming that $g \in H_{n-1}(F)$, we have $y_1 \in F^\times$ and hence we may substitute $b_1 \mapsto b_1 y_1$. Then for $\phi'_{n-1}(b_1(c_{n-2}, 1))$ to be nonzero, we must have $b_1 \in 1 + \varpi^m \mathcal{O}_F$. Note that $\widehat{\phi}_{n-2}$ and φ_{n-2} are invariant under multiplication by $1 + \varpi^m \mathcal{O}_F$. We see that $w(g)$ is given by the product of

$$\eta_n(y_1^{n-1}) \int_{M_{n-1,1}(F)} \phi'_{n-1}(b_1(c_{n-2}, 1)) dc_{n-2} \eta(b_1^{n-1}) d^* b_1, \\ |x_{n-2}|_E \widehat{\phi}_{n-2}(-e_{n-2}^* \tau x_{n-2})$$

and

$$\int_{N_{n-2}(F) \setminus H_{n-2}(F)} \int_{N_{n-2}(E)} \varphi_{n-2}(x_{n-2}^{-1} u_{n-2} a_{n-2}) \overline{\psi}_\tau(u_{n-2}) |a_{n-2}|^{-1} \eta_n(a_{n-2}) du_{n-2} da_{n-2}.$$

When $y = y_1 \cdot \text{diag}(x_{n-2}, 1)$, $x_{n-2} \in B_{n-2,-}(F)$, we have

$$\delta_{n-1}(y) = \delta_{n-2}(x_{n-2}) \det(x_{n-2}).$$

Note that

$$|x_{n-2}|_E = |x_{n-2}|_F^2.$$

Now we may repeat this process to complete the proof. $\square$

For admissible $\varphi_{n-1} \otimes \phi_{n-1} \in \mathcal{C}_c^\infty(H_n(E) \times M_{n,1}(E))$, the function $\widetilde{W}_{\varphi_{n-1}, \phi_{n-1}} \in \mathcal{W}$ lies in $\mathcal{C}_c^\infty(N_{n-1}(E) \setminus H_{n-1}(E), \psi_E)$ and therefore determines a unique element denoted by $W_{\varphi_{n-1}, \phi_{n-1}}$ in $\mathcal{W}$ characterized by

$$(7.5) \quad W_{\varphi_{n-1}, \phi_{n-1}} \begin{pmatrix} g & 0 \\ 0 & 1 \end{pmatrix} = \widetilde{W}_{\varphi_{n-1}, \phi_{n-1}}(g), \quad g \in H_{n-1}(E).$$

Smoothing local Flicker–Rallis period β_n . Let Π_n be an irreducible unitary generic representation of $H_n(E)$. We now consider the local period β_n . We let $\mathcal{W} = \mathcal{W}(\Pi_n, \overline{\psi_E})$ be the Whittaker model of Π_n with respect to the complex conjugate of ψ_E for later convenience. As earlier we have endowed it with a non-degenerate positive definite invariant Hermitian structure:

$$\langle W, W' \rangle = \vartheta(W, W') = \int_{N_{n-1}(E) \backslash H_{n-1}(E)} W \begin{pmatrix} g & \\ & 1 \end{pmatrix} \overline{W' \begin{pmatrix} g & \\ & 1 \end{pmatrix}} dg.$$

We also consider its Kirillov model denoted by $\mathcal{K} = \mathcal{K}(\Pi_n, \overline{\psi_E})$, which is a certain subspace of smooth functions $\mathcal{C}^\infty(N_{n-1}(E) \backslash H_{n-1}(E), \overline{\psi_E})$. Moreover, the Kirillov model always contains the subspace $\mathcal{C}_c^\infty(N_{n-1}(E) \backslash H_{n-1}(E), \overline{\psi_E})$ of smooth compactly-supported functions.

Let $\mathcal{W}^*$ be the (conjugate) algebraic dual space of $\mathcal{W}$ and $\mathcal{H}$ be the Hilbert space underlying the unitary representation Π_n . Then $\mathcal{W}$ is the space of smooth vectors in $\mathcal{H}$ and we have inclusions:

$$\mathcal{W} \subset \mathcal{H} \subset \mathcal{W}^*.$$

The Hermitian pairing on $\mathcal{W} \times \mathcal{W}$ extends to $\mathcal{H} \times \mathcal{H}$ and $\mathcal{W} \times \mathcal{W}^*$. We still denote by Π_n the representation of $H_n(E)$ on $\mathcal{W}^*$ so for any $W \in \mathcal{W}, W' \in \mathcal{W}^*$:

$$\langle \Pi_n(g)W, W' \rangle = \langle W, \Pi_n(g^{-1})W' \rangle.$$

Then the local Flicker–Rallis period β_n is an element in $\mathcal{W}^*$ defined by the convergent integral

$$(7.6) \quad \beta_n(W) = \int_{N_{n-1}(F) \backslash H_{n-1}(F)} W \begin{pmatrix} \epsilon_{n-1}h & \\ & 1 \end{pmatrix} \eta_n(h) dh.$$

To ease notation, we write this as

$$\beta_n(W) = \int_{N_{n-1}(F) \backslash H_{n-1}(F)} W(\epsilon_{n-1}h) \eta_n(h) dh.$$

We also write this as

$$\beta_n(W) = \langle W, \beta_n \rangle.$$

It is $(H_n(F), \eta_n)$ -invariant ([9]):

$$\beta_n \in \text{Hom}_{H_n(F)}(\mathcal{W}, \mathbb{C}_{\eta_n}).$$

We would like to smoothen the local period β_n by applying some sort of “mollifier”.

Proposition 7.2. *Assume that the pair $\varphi_{n-1} \in \mathcal{C}_c^\infty(H_n(E)), \phi_{n-1} \in \mathcal{C}_c^\infty(M_{n,1}(E))$ is m -admissible for $m > 0$. Let $W_{\varphi_{n-1}, \phi_{n-1}} \in \mathcal{W}$ be the element determined by 7.5. Then for every $W \in \mathcal{W}(\Pi_n, \overline{\psi_E})$, we have*

$$\langle \Pi_n(\varphi_{n-1}^*) \Pi_n(\phi_{n-1}^*) W, \beta_n \rangle = \langle W, W_{\varphi_{n-1}, \phi_{n-1}} \rangle.$$

In other words, the linear functional $\Pi_n(\phi_{n-1}) \Pi_n(\varphi_{n-1}) \beta_n$, a priori only in $\mathcal{W}^*$, is indeed a smooth vector and is represented by $W_{\varphi_{n-1}, \phi_{n-1}} \in \mathcal{W}(\Pi_n, \overline{\psi_E})$.

Proof. The LHS is given by

$$\int_{N_{n-1}(F) \backslash H_{n-1}(F)} \int_{H_{n-1}(E)} \Pi_n(\phi_{n-1}^*) W(\epsilon_{n-1} h g_{n-1}) \overline{\varphi}_{n-1}(g_{n-1}^{-1}) dg_{n-1} \eta_n(h) dh.$$

Substitute $g_{n-1} \mapsto h^{-1} \epsilon_{n-1}^{-1} g_{n-1}$,

$$\int_{N_{n-1}(F) \backslash H_{n-1}(F)} \int_{H_{n-1}(E)} \Pi_n(\phi_{n-1}^*) W(g_{n-1}) \overline{\varphi}_{n-1}(g_{n-1}^{-1} \epsilon_{n-1} h) dg_{n-1} \eta_n(h) dh.$$

Since $W(ug_{n-1}) = \overline{\psi}_E(u) W(g_{n-1})$ for $u \in N_{n-1}(E)$, we may rewrite the integral as

$$\int \Pi_n(\phi_{n-1}^*) W(g_{n-1}) \left(\int_{N_{n-1}(E)} \overline{\varphi}_{n-1}(g_{n-1}^{-1} u^{-1} \epsilon_{n-1} h) \overline{\psi}_E(u) du \right) \eta_n(h) dh dg_{n-1},$$

where the outer integral is over

$$h \in N_{n-1}(F) \backslash H_{n-1}(F), \quad g_{n-1} \in N_{n-1}(E) \backslash H_{n-1}(E).$$

Now we also note that

$$\begin{aligned} \Pi_n(\phi_{n-1}^*) W(g_{n-1}) &= \int_{M_{n-1,1}(E)} W \left[g_{n-1} \begin{pmatrix} 1_{n-1} & u \\ & 1 \end{pmatrix} \right] \phi_{n-1}^*(u) du \\ &= W(g_{n-1}) \int_{M_{n-1,1}(E)} \overline{\varphi}_{n-1}(-u) \overline{\psi}_E(e_{n-1}^* g_{n-1} u) du \\ &= W(g_{n-1}) \overline{\phi}_{n-1}(-e_{n-1}^* g_{n-1}). \end{aligned}$$

This completes the proof. $\square$

Smoothing local Rankin-Selberg period λ . Now let $\Pi = \Pi_n \otimes \Pi_{n+1}$ be an irreducible unitary generic representation of $G'(F) = H_n(E) \times H_{n+1}(E)$. We now need another compactness lemma. For an integer $m' > 0$, let $\mathcal{C}_c^\infty(M_{n,1}(E))_{m'}^\spadesuit$ be the subspace consisting of $\phi_n \in \mathcal{C}_c^\infty(M_{n,1}(E))$ such that the Fourier transform $\hat{\phi}$ is supported in the domain

$$\{(x_1, \dots, x_n) \in M_{n,1}(E) \mid |x_i/x_n| \leq |\varpi_E|^{m'}, i = 1, 2, \dots, n-1.\}$$

Lemma 7.3. *Let $\phi_n \in \mathcal{C}_c^\infty(M_{n,1}(E))_{m'}^\spadesuit$. Let $W \in \mathcal{W}(\Pi_n, \overline{\psi}_E)$ such that its restriction to $H_{n-1}(E)$ has support compact modulo $N_{n-1}(E)$. If m' is sufficiently large (depending on the vector W), the map*

$$H_n(E) \ni g \mapsto \hat{\phi}_n(e_n^* g) W(g)$$

defines an element in $\mathcal{C}_c^\infty(N_n(E) \backslash H_n(E), \overline{\psi}_E)$.

Proof. Clearly $\widehat{\phi}_n(e_n^*g)$ is $N_n(E)$ -invariant and smooth hence the product defines an element in $\mathcal{C}^\infty(N_n(E)\backslash H_n(E), \overline{\psi}_E)$. It remains to show that the product has support compact modulo $N_n(E)$. By the support condition of $\widehat{\phi}_n(e_n^*g)$, we may assume that the lower right entry of g is non-zero. Therefore we may write $g = xu \begin{pmatrix} h & \\ & 1 \end{pmatrix} \begin{pmatrix} 1_{n-1} & \\ z & 1 \end{pmatrix}$ where $h \in H_{n-1}(E), u \in N_n(E), z \in M_{1,n-1}(E)$ and $x \in E^\times$. By the assumption on $\widehat{\phi}_n$, we may assume that $\|z\| < |\varpi_E|^{m'}$ (otherwise the product vanishes). We may choose m' sufficiently large so that W is invariant under right multiplication by $1 + \varpi_E^{m'} M_{n,n}(\mathcal{O}_E)$ (such m' exists since $W \in \mathcal{W}$). We now have

$$\begin{aligned} \widehat{\phi}_n(e_n^*g)W(g) &= \psi_E(u)\widehat{\phi}_n(x(z, 1))W \left[x \begin{pmatrix} h & \\ & 1 \end{pmatrix} \right] \\ &= \psi_E(u)\omega_{\Pi_n}(x)\widehat{\phi}_n(x(z, 1))W \left[\begin{pmatrix} h & \\ & 1 \end{pmatrix} \right], \end{aligned}$$

where ω_{Π_n} is the central character of Π_n . Since the last factor has support compact modulo $N_{n-1}(E)$, and x lies in a compact region, the proof is complete. $\square$

Since the Kirillov model $\mathcal{K}(\Pi_{n+1}, \overline{\psi}_E)$ contains $\mathcal{C}_c^\infty(N_n(E)\backslash H_n(E), \overline{\psi}_E)$ as a subspace, we may view the product $\widehat{\phi}_n(e_n^*g)W(g)$ as an element in $\mathcal{K}(\Pi_{n+1}, \overline{\psi}_E)$. This also determines uniquely an element denoted by W_{ϕ_n} in the Whittaker model $\mathcal{W}(\Pi_{n+1}, \overline{\psi}_E)$. In other words, to each W whose restriction to $H_{n-1}(E)$ lies in $\mathcal{C}_c^\infty(N_{n-1}(E)\backslash H_{n-1}(E), \overline{\psi}_E)$ and a $\phi_n \in \mathcal{C}_c^\infty(M_{n,1}(E))_{m'}^\dagger$ for m' sufficiently large, we associate an element $W_{\phi_n} \in \mathcal{W}(\Pi_{n+1}, \overline{\psi}_E)$ characterized by

$$(7.7) \quad W_{\phi_n} \begin{pmatrix} g & \\ & 1 \end{pmatrix} = \widehat{\phi}_n(e_n^*g)W(g), \quad g \in H_n(E).$$

Its complex conjugate $\overline{W}_{\phi_n}$ defines an element in $\mathcal{W}(\Pi_{n+1}, \psi_E)$.

Now recall that the local Rankin-Selberg period is defined by the convergent integral, cf. (3.13):

$$\lambda(W \otimes W') = \int_{N_n(E)\backslash H_n(E)} W(g)W' \begin{pmatrix} g & \\ & 1 \end{pmatrix} dg,$$

where $W \in \mathcal{W}(\Pi_n, \overline{\psi}_E), W' \in \mathcal{W}(\Pi_{n+1}, \psi_E)$. For $\phi_n \in \mathcal{C}_c^\infty(M_{n,1}(E))$, we have an action on $\mathcal{W}(\Pi_{n+1}, \psi_E)$ by:

$$\Pi_{n+1}(\phi_n)W'(g) = \int_{M_{n-1,1}(E)} W \left[g \begin{pmatrix} 1_{n-1} & u \\ & 1 \end{pmatrix} \right] \phi(u) du, \quad g \in H_{n+1}(E).$$

Proposition 7.4. *Assume that $W \in \mathcal{W}(\Pi_n, \overline{\psi}_E)$ whose restriction to $H_{n-1}(E)$ lies in*

$$\mathcal{C}_c^\infty(N_{n-1}(E)\backslash H_{n-1}(E), \overline{\psi}_E)$$

and that $\hat{\phi}_n \in \mathcal{C}_c^\infty(M_{n,1}(E))_{m'}^\dagger$ for m' sufficiently large. Then for every $W' \in \mathcal{W}(\Pi_{n+1}, \psi_E)$ we have

$$\lambda(W \otimes \Pi_{n+1}(\phi_n)W') = \langle W', \overline{W}_{\phi_n} \rangle.$$

Proof. For $g \in H_n(E)$, we have

$$\begin{aligned} \Pi_{n+1}(\phi_n)W' \begin{pmatrix} g & \\ & 1 \end{pmatrix} &= \int_{M_{n-1,1}(E)} W' \left[\begin{pmatrix} g & \\ & 1 \end{pmatrix} \begin{pmatrix} 1_{n-1} & u \\ & 1 \end{pmatrix} \right] \phi_n(u) du \\ &= \int_{M_{n-1,1}(E)} W' \left[\begin{pmatrix} g & \\ & 1 \end{pmatrix} \right] \psi_E(e_n^* g u) \phi(u) du \\ &= \hat{\phi}_n(e_n^* g) W' \left[\begin{pmatrix} g & \\ & 1 \end{pmatrix} \right]. \end{aligned}$$

Return to the local Rankin-Selberg period:

$$\begin{aligned} \lambda(W \otimes \Pi_{n+1}(\phi_n)W') &= \int_{N_n(E) \backslash H_n(E)} W(g) \Pi_{n+1}(\phi_n)W' \begin{pmatrix} g & \\ & 1 \end{pmatrix} dg \\ &= \int_{N_n(E) \backslash H_n(E)} W(g) \hat{\phi}_n(e_n^* g) W' \begin{pmatrix} g & \\ & 1 \end{pmatrix} dg \\ &= \langle W', \overline{W}_{\phi_n} \rangle. \end{aligned}$$

This completes the proof. □

8 Local character expansion in the general linear case

In this section we prove a “limit” formula for the (local) spherical character in the general linear side. We will choose a subspace of test functions supported in a small neighborhood of the origin of the symmetric space S_{n+1} . Then we may treat them as functions on the “Lie algebra” $\mathfrak{s}_{n+1}$ of S_{n+1} , the tangent space at origin. The key property for these functions is that their Fourier transform have vanishing unipotent orbital integrals except for the regular unipotent element. The intermediate steps are messy and somehow ugly; but the final outcome seems to be miraculously neat.

A “limit” formula for the spherical character $I_\Pi(f)$. We now combine the two propositions above to obtain a formula of $I_\Pi(f)$ for a special class of test functions f . Now let $\Pi = \Pi_n \otimes \Pi_{n+1}$ where the central character of both Π_n and Π_{n+1} are trivial on $F^\times$.

Consider $f_n \in \mathcal{C}_c^\infty(H_n(E))$ and $f_{n+1} \in \mathcal{C}_c^\infty(H_{n+1}(E))$. Let $\phi_n \in \mathcal{C}_c^\infty(M_{n,1}(E))$. We consider a *perturbation* of f_{n+1} by ϕ_n

$$f_{n+1}^{\phi_n}(g) = \int_{M_{n,1}(E)} f_{n+1} \left[\begin{pmatrix} 1_n & u \\ & 1 \end{pmatrix} g \right] \phi(u) du, \quad g \in H_{n+1}(E).$$

This is the same as $\phi_n \circ f_{n+1}$ —the convolution introduced in §7—where we view $M_{n,1}(E)$ as a subgroup of $H_{n+1}(E)$. Similarly, for $\varphi_{n-1} \in \mathcal{C}_c^\infty(H_{n-1}(E))$, $\phi_{n-1} \in \mathcal{C}_c^\infty(M_{n-1,1}(E))$, we define a *perturbation* of f_n : for $g \in H_n(E)$,

$$f_n^{\varphi_{n-1}, \phi_{n-1}}(g) := \int_{M_{n,1}(E)} \int_{H_{n-1}(E)} f_2 \left[g \begin{pmatrix} x^{-1} & \\ & 1 \end{pmatrix} \begin{pmatrix} 1 & -u \\ & 1 \end{pmatrix} \right] \varphi_{n-1}(x) \phi_{n-1}(u) dx du.$$

Equivalently

$$f_n^{\varphi_{n-1}, \phi_{n-1}} = f_n \circ \phi_{n-1} \circ \varphi_{n-1} \in \mathcal{C}_c^\infty(H_n(E)).$$

We will consider functions of the following form:

$$f = f_n^{\varphi_{n-1}, \phi_{n-1}} \otimes f_{n+1}^{\phi_n} \in \mathcal{C}_c^\infty(H_n(E) \times H_{n+1}(E)).$$

Definition 8.1. Fix Π . Let (m, m', r) be positive integers with $r > m' > m > 0$. We say that $f = f_n^{\varphi_{n-1}, \phi_{n-1}} \otimes f_{n+1}^{\phi_n}$ is (m, m', r) -admissible or admissible for Π if it satisfies the following:

- The function $\varphi_{n-1} \otimes \phi_{n-1}$ is m -admissible. Hence it determines an element $\widetilde{W}_{\varphi_{n-1}, \overline{\phi}_{n-1}} \in \mathcal{C}_c^\infty(N_{n-1} \backslash H_{n-1}(E), \psi_E)$ (cf. Lemma 7.1) and $W_{\varphi_{n-1}, \phi_{n-1}} \in \mathcal{W}(\Pi_n, \psi_E)$.
- The function $\phi_n \in \mathcal{C}_c^\infty(M_{n,1}(E))_{m'}^\spadesuit$ for sufficiently large m' (depending on Π_n , $\varphi_{n-1} \otimes \phi_{n-1}$ and hence on the integer m). More precisely, m' is large enough such that Lemma 7.3 holds for $W = W_{\varphi_{n-1}, \phi_{n-1}}$. Let $W_{\varphi_{n-1}, \phi_{n-1}, \phi_n} \in \mathcal{W}(\Pi_{n+1}, \overline{\psi}_E)$ be the function characterized by the equation (7.7) for the choice $W = W_{\varphi_{n-1}, \phi_{n-1}}$.
- The function f_n (f_{n+1} , resp.) is a multiple of the characteristic function of $1 + \varpi^r M_{n,n}(\mathcal{O}_E)$ (of $1 + \varpi^r M_{n+1,n+1}(\mathcal{O}_E)$, resp.). We normalize f_n, f_{n+1} by

$$(8.1) \quad \int_{H_n(E)} f_n(g) dg = \int_{H_{n+1}(E)} f_{n+1}(g) dg = 1.$$

We require that r is sufficiently large (depending on Π , $\varphi_{n-1}, \phi_{n-1}, \phi_n$ and hence on m, m') so that

$$\Pi_n(f_n) W_{\varphi_{n-1}, \phi_{n-1}} = W_{\varphi_{n-1}, \phi_{n-1}}$$

and

$$\Pi_{n+1}(f_{n+1}) W_{\varphi_{n-1}, \phi_{n-1}, \phi_n} = W_{\varphi_{n-1}, \phi_{n-1}, \phi_n}.$$

Remark 8. It will be useful later to note that if $f = f_n^{\varphi_{n-1}, \phi_{n-1}} \otimes f_{n+1}^{\phi_n}$ is (m, m', r) -admissible, then $f_{t,t'}$ is (m) -admissible.

Proposition 8.2. Fix Π and assume that $f = f_n^{\varphi_{n-1}, \phi_{n-1}} \otimes f_{n+1}^{\phi_n}$ is (m, m', r) -admissible for Π . Then we have

$$I_\Pi(f) = \beta_{n+1}(W_{\varphi_{n-1}, \phi_{n-1}, \phi_n}).$$

Proof. First we have

$$\begin{aligned} I_{\Pi}(f_n \otimes f_{n+1}) &= \sum_{W, W'} \lambda(\Pi_n(f_n)W \otimes \Pi_{n+1}(f_{n+1})W') \overline{\beta_n(W)} \cdot \overline{\beta_{n+1}(W')} \\ &= \sum_{W, W'} \lambda(W \otimes \Pi_{n+1}(f_{n+1})W') \overline{\beta_n(\Pi_n(f_n^*)W)} \cdot \overline{\beta_{n+1}(W')}, \end{aligned}$$

where the sum of W (W' , resp.) runs over an orthonormal basis of Π_n (Π_{n+1} , resp.).

For simplicity, we write ϕ for ϕ_{n-1} and φ for φ_{n-1} . Now we replace f_n by $f_n^{\varphi, \phi} = f \circ \phi \circ \varphi$. Note that

$$\Pi_n((f_n^{\varphi, \phi})^*) = \Pi_n(\varphi^*)\Pi_n(\phi^*)\Pi_n(f^*)$$

By Prop. 7.2, we have for all $W \in \mathcal{W}(\Pi_n, \overline{\psi_E})$

$$\beta_n(\Pi_n((f_n^{\varphi, \phi})^*)W) = \langle \Pi_n(f_n^*)W, W_{\varphi, \phi} \rangle = \langle W, \Pi_n(f_n)W_{\varphi, \phi} \rangle.$$

For any $W_0 \in \mathcal{W}(\Pi_n, \overline{\psi_E})$, we have an orthonormal expansion

$$W_0 = \sum_W \langle W_0, W \rangle W,$$

where the sum of W runs over an orthonormal basis of Π_n . Hence we may fold the sum over W to obtain

$$I_{\Pi}(f_n^{\varphi, \phi} \otimes f_{n+1}) = \sum_{W'} \lambda(\Pi_n(f_n)W_{\varphi, \phi} \otimes \Pi_{n+1}(f_{n+1})W') \overline{\beta_{n+1}(W')}.$$

Now we further replace f_{n+1} by $f_{n+1}^{\phi_n} = \phi_n \circ f_{n+1}$ and assume that $f = f_n^{\varphi, \phi} \otimes f_{n+1}^{\phi_n}$ is admissible. By the admissibility, we have

$$\Pi_n(f_n)W_{\varphi, \phi} = W_{\varphi, \phi}.$$

and

$$\Pi_{n+1}(f_{n+1}^*)\overline{W}_{\varphi, \phi, \phi_n} = \overline{W}_{\varphi, \phi, \phi_n}.$$

By Prop. 7.4 and $\int_{H_n(E)} f_n(g)dg = 1$, we have for all $W' \in \mathcal{W}(\Pi_{n+1}, \psi_E)$:

$$\begin{aligned} \lambda(\Pi_n(f_n)W_{\varphi, \phi} \otimes \Pi_{n+1}(f_{n+1}^{\phi_n})W') &= \langle \Pi_{n+1}(f_{n+1})W', \overline{W}_{\varphi, \phi, \phi_n} \rangle \\ &= \langle W', \Pi_{n+1}(f_{n+1}^*)\overline{W}_{\varphi, \phi, \phi_n} \rangle \\ &= \langle W', \overline{W}_{\varphi, \phi, \phi_n} \rangle. \end{aligned}$$

Then we may fold the sum over W' to obtain

$$I_{\Pi}(f_n^{\varphi, \phi} \otimes f_{n+1}^{\phi_n}) = \overline{\beta_{n+1}(\overline{W}_{\varphi, \phi, \phi_n})} = \beta_{n+1}(W_{\varphi, \phi, \phi_n}).$$

This completes the proof. □

We need to simplify the formula. Define for $a \in A_n(F)$:

$$(8.2) \quad \bar{\delta}_n(a) := \det(\text{Ad}(a) : \mathfrak{n}_n) / \det(\text{Ad}(a) : \mathfrak{n}_{n-1}).$$

To simplify the exposition from now on we redefine $f_{n+1}^{\phi_n}$ as $\phi_n^\vee \circ f_{n+1}$.

Proposition 8.3. Fix Π and assume that $f = f_n^{\varphi_{n-1}, \phi_{n-1}} \otimes f_{n+1}^{\phi_n}$ is (m, m, r) -admissible for Π . Use notations as in Lemma 7.1. Then we have

$$I_\Pi(f) = \omega_{\Pi_n}(\tau) |\tau|_E^{d_n} \left(\int_{B_{n-1,-}(F)} \phi'(b) db \right) \int \prod_{i=0}^{n-1} \widehat{\phi_{n-i}}(-y_i(z_{n-i-1}, 1)\tau) |\bar{\delta}_n(y')|^{-1} \eta(y') d^*y' dz,$$

where the integral of $d^*y' dz$ is over $y' \in A_n(F)$, $z \in N_{n,-}(F)$:

$$y' = \begin{pmatrix} y_0 y_1 y_2 \dots y_{n-1} & & \\ & \ddots & \\ & & y_0 y_1 & \\ & & & y_0 \end{pmatrix} \in A_n(F),$$

and

$$z' = \prod_{i=1}^{n-1} \begin{pmatrix} 1_i & \\ z_i & 1 \end{pmatrix} \in N_{n,-}(F), \quad z_i \in M_{1,i}(F).$$

Remark 9. Note that $|\tau|_E^{d_n} = |\det(\text{Ad}(\epsilon_{n-1}) : N_{n-1}(E))|$. If we consider the normalized distribution

$$|\tau|_E^{-(d_n + d_{n+1})/2} I_\Pi(f) = |\det(\text{Ad}(\epsilon_{n-1}) : N_{n-1}(E))|_E^{-1/2} |\det(\text{Ad}(\epsilon_n) : N_n(E))|_E^{-1/2} I_\Pi(f),$$

then it is independent of the choice of the generator τ (up to a sign).

Proof. By (7.7), we have (cf. (7.6), and note to replace ϕ_n by $\phi_n^\vee$)

$$\beta_{n+1}(W_{\varphi, \phi, \phi_n}) = \int_{N_n(F) \backslash H_n(F)} W_{\varphi, \phi}(\epsilon_n h) \hat{\phi}_n(-e_n^* \epsilon_n h) \eta_{n+1}(h) dh.$$

Let $h = y' z'$ as in the Lemma and

$$y = y_0 \begin{pmatrix} y & \\ & 1 \end{pmatrix} \in A_n(F), y \in A_{n-1}(F),$$

and

$$z' = \begin{pmatrix} z & \\ & 1 \end{pmatrix} \begin{pmatrix} 1_{n-1} & \\ z_{n-1} & 1 \end{pmatrix}, z \in N_{n-1,-}(F).$$

Then we may replace the integral on the quotient $N_n(F) \backslash H_n(F)$ by the integral over y', z' for the measure:

$$|\delta_n(y')|^{-1} \prod_{i=0}^{n-1} d^* y_i \prod_{j=1}^{n-1} dz_j.$$

Since $\delta_n(y') = \delta_{n-1}(y) \det(y) = \delta_{n-1}(y) \bar{\delta}_n(y')$, we may also write this as a product

$$(|\delta_{n-1}(y)|^{-1} |y|^{-1} d^* y dz) d^* y_0 dz_{n-1},$$

where $d^* y = \prod_{i=0}^{n-2} d^* y_i$ and $dz = \prod_{j=1}^{n-2} dz_j$. Since the central character of Π_n is trivial on $F^\times$, we have

$$W_{\varphi, \phi}(\epsilon_n y_0 h) = W_{\varphi, \phi}(\epsilon_n h).$$

By the admissibility, the value $\hat{\phi}_n(-e_n^* \epsilon_n h) = \hat{\phi}_n(-y_0(z_{n-1}, 1)\tau)$ is non-zero only if $\|z_{n-1}\|$ is very small so that $\begin{pmatrix} 1_{n-1} & \\ z_{n-1} & 1 \end{pmatrix}$ acts trivially on $W_{\varphi, \phi}$. This allows us to write the integral as the product of

$$\int \hat{\phi}_n(-y_0(z_{n-1}, 1)\tau) \eta_{n+1}(y_0^n) d^* y_0 dz_{n-1}$$

and

$$\omega_{\Pi_n}(\tau) \int_{A_{n-1}(F)} \int_{N_{n-1,-}(F)} W_{\varphi, \phi} \left(\begin{pmatrix} \epsilon_{n-1} y z & \\ & 1 \end{pmatrix} \right) |\delta_{n-1}(y)|^{-1} |y|^{-1} \eta_{n+1}(y) d^* y dz,$$

where we have used the equality

$$\epsilon_n = \tau \begin{pmatrix} \epsilon_{n-1} & \\ & 1 \end{pmatrix}.$$

By definition $W_{\varphi, \phi} \left(\begin{pmatrix} \epsilon_{n-1} y z & \\ & 1 \end{pmatrix} \right) = \widetilde{W}_{\varphi, \phi}(\epsilon_{n-1} y z)$ (cf. (7.5)), we may apply the formula of the latter by Lemma 7.1:

$$|\tau|_E^{d_n} \left(\int_{B_{n-1,-}(F)} \phi'(b) db \right) \eta_n(y) |\delta_{n-1}(y)|_F \prod_{i=1}^{n-1} \widehat{\phi_{n-i}}(y_i(z_{n-i-1}, 1)\tau).$$

Finally we note $\eta_n \eta_{n+1} = \eta$ and $\eta_{n+1}(y_0^n) = \eta(y_0^{n^2}) = \eta(y_0^n)$. This completes the proof. $\square$

Truncated local expansion of the spherical character I_Π . We are now ready to deduce a truncated local expansion of I_Π around the origin. As we have alluded in the Introduction, this expansion is the relative version of Harish-Chandra's local expansion for characters of admissible representations of p -adic reductive groups in terms of Fourier transform of nilpotent orbital integrals. Here we obtain a truncated expansion that only involves the regular unipotent element.

First we associate to $f \in \mathcal{C}_c^\infty(H_n(E) \times H_{n+1}(E))$ with small support around 1 a function on the Lie algebra $\mathfrak{s}$ with small support around 0. To $f = f_n \otimes f_{n+1} \in \mathcal{C}_c^\infty(H_n(E) \times H_{n+1}(E))$ we associate a function $\tilde{f} \in \mathcal{C}_c^\infty(H_{n+1}(E))$ by

$$\tilde{f}(g) = \int_{H_n(E)} f_n(h) f_{n+1}(hg) dh.$$

In other words, we have $\tilde{f} = f_n^\vee \circ f_{n+1}$. Moreover, we also have

$$\begin{aligned} \nu &= \nu_{n+1} : H_{n+1,E} \rightarrow S_{n+1} \\ g &\mapsto g\bar{g}^{-1}. \end{aligned}$$

The fiber integral defines a map denoted by $\nu : \mathcal{C}_c^\infty(H_{n+1}(E)) \rightarrow \mathcal{C}_c^\infty(S_{n+1}(F))$:

$$\nu(\tilde{f})(x) := \int_{H_{n+1}(F)} \tilde{f}(gh) dh, \quad \nu(g) = x,$$

if n is even, and

$$\nu(\tilde{f})(x) := \int_{H_{n+1}(F)} \tilde{f}(gh) \eta'(gh) dh, \quad \nu(g) = x,$$

when n is odd (then this depends on the auxiliary character η'). The Cayley map is a local isomorphism near a neighborhood of $0 \in \mathfrak{s}$

$$\begin{aligned} \mathfrak{c} &= \mathfrak{c}_{n+1} : \mathfrak{s} \rightarrow S_{n+1} \\ X &\mapsto \frac{1+X}{1+\overline{X}} = \frac{1+X}{1-X}. \end{aligned}$$

and its inverse is given by

$$\mathfrak{c}^{-1}(x) = -\frac{1-x}{1+x}.$$

In particular, for a function $\Phi \in \mathcal{C}_c^\infty(S_n)$ ($\phi \in \mathcal{C}_c^\infty(S_n)$, resp.) with support in a small neighborhood of $1 \in S_{n+1}$ ($0 \in \mathfrak{s}$, resp.), we may consider it as a function on $\mathfrak{s}$ (S_{n+1} , resp.) denoted by $\mathfrak{c}^{-1}(\Phi)$ ($\mathfrak{c}(\phi)$), resp.). We also have a morphism:

$$\begin{aligned} \iota &= \iota_{n+1} : \mathfrak{s} \rightarrow H_{n+1,E} \\ X &\mapsto 1+X, \end{aligned}$$

such that, wherever they are all defined, we have $\nu \circ \iota = \mathfrak{c}$:

$$\begin{array}{ccc} & & H_{n+1}(E) \\ & \nearrow \iota & \downarrow \nu \\ \mathfrak{s}_{n+1} & \xrightarrow{\mathfrak{c}} & S_{n+1} \end{array}$$

We associate to $f \in \mathcal{C}_c^\infty(H_n(E) \times H_{n+1}(E))$ a function on $\mathfrak{s}$ denoted by $f_{\mathfrak{h}}$

$$(8.3) \quad f_{\mathfrak{h}}(X) := \int_{H_n(F)} \tilde{f}((1+X)h)dh, \quad X \in \mathfrak{s}.$$

Then we have when $\det(1-X) \neq 0$

$$\mathfrak{c}^{-1}(\nu(\tilde{f}))(X) = f_{\mathfrak{h}}(X), \quad X \in \mathfrak{s}.$$

From now on all of test functions will be supported in small neighbourhoods of the obvious distinguished points of $H_n(E) \times H_{n+1}(E)$, $S_{n+1}(F)$ and $\mathfrak{s}(F)$. In particular, we have $f_{\mathfrak{h}} \in \mathcal{C}_c^\infty(\mathfrak{s})$.

We then consider $\mathfrak{s}(F)$ as a subspace of $M_{n+1,n+1}(E)$. On $M_{n+1,n+1}(E)$ we have a bilinear pairing $\langle X, Y \rangle := \text{tr}(XY)$, under which the decomposition $M_{n+1,n+1}(E) = M_{n+1,n+1}(F) \oplus \mathfrak{s}(F)$ is orthogonal. We then define the Fourier transform on $\mathfrak{s}(F)$ with respect to the restriction of the above pairing:

$$\hat{f}_{\mathfrak{h}}(X) := \int_{\mathfrak{s}} f_{\mathfrak{h}}(Y) \psi(\text{tr}(XY)) dY,$$

where the measure is then the self-dual measure for the Fourier transform. We will consider the orbital integral of the regular unipotent element

$$\xi_- = \xi_{n+1,-} = \tau \begin{pmatrix} 0 & 0 & \dots & 0 \\ 1 & 0 & 0 & \dots \\ \dots & 1 & 0 & 0 \\ 0 & \dots & 1 & 0 \end{pmatrix} \in \mathfrak{s}(F).$$

Theorem 8.4. *Let Π be a unitary generic irreducible admissible representation. Then for sufficiently admissible $f \in \mathcal{C}_c^\infty(H_n(E) \times H_{n+1}(E))$, we have*

$$I_{\Pi}(f) = |\tau|_E^{(d_n+d_{n+1})/2} \omega_n(\tau) \cdot \mu_{\xi_-}(\hat{f}_{\mathfrak{h}}),$$

where $\omega_n = \omega_{\Pi_n}$ is the central character of Π_n .

Remark 10. Note that $\mu_{\xi_-}(\hat{f}_{\mathfrak{h}})$ is independent of the choice of τ (up to a sign).

The proof will occupy the rest of this section.

Determine $f_{\mathfrak{h}}$. We consider admissible functions of the form $f_n^{\varphi_{n-1}, \phi_{n-1}} \otimes f_{n+1}^{\phi_n^\vee}$ —note the change to $\phi_n^\vee$ to ease later exposition. Let P —not to be confused with the mirabolic P_n —be the subgroup P of $H_{n+1}(E)$ consisting of elements

$$\begin{pmatrix} * & * & * & * \\ * & \dots & * & * \\ 0 & 0 & 1 & * \\ 0 & 0 & 0 & 1 \end{pmatrix} \in H_{n+1}(E) \subset M_{n+1,n+1}(E).$$

The triple $(\varphi_{n-1}, \phi_{n-1}, \phi_n)$ then defines a function on Ψ by

$$\Psi(u'ux) := \varphi_{n-1}(x)\phi_{n-1}(u)\phi_n(u').$$

For simplicity, we write

$$f^\Psi = f_n^{\varphi_{n-1}, \phi_{n-1}} \otimes f_{n+1}^{\phi_n^\vee}.$$

We also consider the Lie algebra $\mathfrak{p}$ of P :

$$\begin{pmatrix} * & * & * & * \\ * & \dots & * & * \\ 0 & 0 & 0 & * \\ 0 & 0 & 0 & 0 \end{pmatrix} \in M_{n+1, n+1}(E).$$

The map $p \mapsto 1 + p$ from $\mathfrak{p}$ to P defines a local homeomorphism from $\varpi\mathfrak{p}(\mathcal{O}_E)$ to its image. We decompose $\mathfrak{p} = \mathfrak{p}^+ \oplus \mathfrak{p}^-$. Then the function Ψ is supported in the subgroup of $P(E)$:

$$P(\varpi\mathcal{O}) \oplus \mathfrak{p}(\varpi\mathcal{O}_E^-) = 1 + \mathfrak{p}(\varpi\mathcal{O}_E).$$

Note that this is a unimodular subgroup. So either the left or right invariant measure on $P(E)$ restricts to a Haar measure on it. Let $(\varphi_{n-1}, \phi_{n-1}, \phi_n)$ be (m, m') -admissible. We may write $\phi_i = \phi^+ \otimes \phi_i^-$ ($i = n-1, n$) according to the decomposition $M_{i,1}(E) = M_{i,1}(F) \oplus M_{i,1}(E^-)$ and similarly for φ_{n-1} viewed as a function on $M_{n-1, n-1}(E)$. Then we define Ψ^+ and Ψ^- as functions on $1 + \mathfrak{p}^+(\mathcal{O}_F)$ and $\mathfrak{p}^-$ respectively:

$$\Psi^? = \varphi_{n-1}^? \otimes \phi_{n-1}^? \otimes \phi_n^?, \quad ? = \pm.$$

We may even assume that Ψ^+ is a multiple of the characteristic function of the subset of $\mathfrak{p}^+$ consisting of matrices (x_{ij}) where $x_{ij} \equiv \delta_{ij} \pmod{\varpi^m}$ when $j \leq n$ and $x_{i, n+1} \equiv \delta_{i, n+1} \pmod{\varpi^{m'}}$. Then it is elementary to check that $\Psi = \Psi^+ \otimes \Psi^-$ and

$$(8.4) \quad \Psi^-(p_+p_-) = \Psi^-(p_-)$$

whenever $p_+ \in \text{supp}(\Psi^+)$.

Lemma 8.5. *Fix Π and assume that $f^\Psi = f_n^{\varphi_{n-1}, \phi_{n-1}} \otimes f_{n+1}^{\phi_n^\vee}$ is admissible for Π . Then we have*

$$f_{\mathfrak{q}}^\Psi(X) = c(\Psi^+) \int_{\mathfrak{p}^-} f_{\mathfrak{q}}(X + p) \Psi^-(p) dp,$$

where $f_{\mathfrak{q}}$ is the function on $\mathfrak{s}$ associated to $f_n \otimes f_{n+1}$ and

$$c(\Psi^+) = \int_{1 + \mathfrak{p}^+(\varpi\mathcal{O})} \Psi^+(p) dp$$

is a constant.

Proof. We consider the case when n is even; the case n even is similar and only require us to change notations several places. By definition we have

$$\widetilde{f}^\Psi = (f_n \circ \phi_{n-1} \circ \varphi_{n-1})^\vee \circ f_{n+1}^{\phi_n^\vee} = \varphi_{n-1}^\vee \circ \phi_{n-1}^\vee \circ f_n^\vee \circ f_{n+1}^{\phi_n^\vee}.$$

Since f_n a multiple of $1_{1+\varpi_E^r M_{n,n}(\mathcal{O}_E)}$ for sufficiently large r and $\int f(g)dg = 1$, we may assume that $f_n^\vee \circ f_{n+1}^{\phi_n^\vee}$ is the same as $f_{n+1}^{\phi_n^\vee}$:

$$\widetilde{f}^\Psi = \varphi_{n-1}^\vee \circ \phi_{n-1}^\vee \circ f_{n+1}^{\phi_n^\vee} = \varphi_{n-1}^\vee \circ \phi_{n-1}^\vee \circ \phi_n^\vee \circ f_{n+1}.$$

Explicitly, this is

$$\widetilde{f}^\Psi(g) = \int \varphi_{n-1}(x) \phi_{n-1}(u) f_{n+1}^{\phi_n^\vee}(uxg) dx du = \int_P \Psi(p) f_{n+1}(pg) dp.$$

Let's denote the RHS by $f_{n+1}^\Psi(g)$. By the admissibility, the function Ψ is of the form $\Psi^+ \otimes \Psi^-$ where $\Psi^\pm \in \mathcal{C}_c^\infty(\mathfrak{p}^\pm)$. By our choice, f_{n+1} is a multiple of $1_{1+\varpi_E^{r'} M_{n+1,n+1}(\mathcal{O}_E)}$ for sufficiently large r' . In particular, f_{n+1} is conjugate invariant under $1 + \varpi_E M_{n+1,n+1}(\mathcal{O}_E)$:

$$f_{n+1}(hgh^{-1}) = f_{n+1}(g), \quad h \in 1 + \varpi_E M_{n+1,n+1}(\mathcal{O}_E).$$

By the support condition, $f_{\mathfrak{h}}^\psi(X)$ vanishes unless $X \in \mathfrak{s}(\mathcal{O}_F)$. We thus assume that $X \in \mathfrak{s}(\mathcal{O}_F)$. Then by definition and the support condition of Ψ :

$$\begin{aligned} f_{\mathfrak{h}}^\Psi(X) &= \int_{H_{n+1}(F)} \int_{P(E)} \Psi(p) f_{n+1}(p(1+X)h) dp dh \\ &= \int_{H_{n+1}(F)} \int_{P(F)} \int_{\mathfrak{p}^-} \Psi(p_+(1+p_-)) f_{n+1}(p_+(1+p_-)(1+X)h) dp_+ dp_- dh. \end{aligned}$$

By the invariance of Ψ^- under the support of Ψ^+ , we have

$$\begin{aligned} &\int_{H_{n+1}(F)} \int_{P(F)} \int_{\mathfrak{p}^-} \Psi^+(p_+) \Psi^-(p_-) f_{n+1}((1+p_-)(1+X)h p_+) dp_+ dp_- dh \\ &= \int_{H_{n+1}(F)} \int_{P(F)} \int_{\mathfrak{p}^-} \Psi^+(p_+) \Psi^-(p_-) f_{n+1}((1+p_-)(1+X)h) dp_+ dp_- dh \\ &= c(\Psi^+) \int_{H_{n+1}(F)} \int_{\mathfrak{p}^-} \Psi^-(p_-) f_{n+1}((1+p_-)(1+X)h) dp_- dh. \end{aligned}$$

Note that

$$(1+p_-)(1+X) = (1+p_-X)(1+(1+p_-X)^{-1}(p_-+X))$$

and $(1+p_-X) \in 1 + \varpi M_{n+1,n+1}(\mathcal{O}_F) \subset H_{n+1}(F)$. We have

$$\begin{aligned} &\int_{H_{n+1}(F)} \int_{\mathfrak{p}^-} \Psi^-(p_-) f_{n+1}((1+p_-)(1+X)h) dp_- dh \\ &= \int_{H_{n+1}(F)} \int_{\mathfrak{p}^-} \Psi^-(p_-) f_{n+1}((1+(1+p_-X)^{-1}(p_-+X))h) dp_- dh. \end{aligned}$$

Compared to the definition of $f_{\mathfrak{h}}$, we obtain

$$f_{\mathfrak{h}}^{\Psi}(X) = c(\Psi^+) \int_{\mathfrak{p}^-} \Psi^-(p_-) f_{\mathfrak{h}}((1 + p_- X)^{-1}(p_- + X)) dp_-.$$

Finally note that $f_{\mathfrak{h}}(X)$ is a multiple of $1_{\varpi^r \mathfrak{s}(\mathcal{O}_F)}$ for some large integer r . It follows that $f_{n+1, \mathfrak{h}}(X) = f_{n+1, \mathfrak{h}}(hX)$ for any $h \in 1 + \varpi M_n(\mathcal{O}_F)$. Since $p_- \in \text{supp}(\Psi^-) \subset \mathfrak{p}^-(\varpi \mathcal{O})$ and $X \in \mathfrak{s}(\mathcal{O})$, we therefore obtain

$$f_{\mathfrak{h}}^{\Psi}(X) = c(\Psi^+) \int_{\mathfrak{p}^-} \Psi^-(p_-) f_{n+1, \mathfrak{h}}(p_- + X) dp_-.$$

□

Local constancy of the orbital integral. To compare with the unitary case in the next section, we need to understand the orbital integral of $\widehat{f_{\mathfrak{h}}^{\Psi}}$, at least around zero. We show that the orbital interval is locally constant around zero and the constant is essentially given by the regular unipotent orbital integral. The result will be used in the next section.

First we study the support of $\widehat{f_{\mathfrak{h}}^{\Psi}}$, especially its restriction to the relative nilpotent cone $\mathcal{N}$.

Lemma 8.6. *Let f be a sufficiently admissible function. Decompose $\mathfrak{s}_{n+1} = \mathfrak{s}_n \oplus \mathfrak{s}_n^{\perp}$ and we may write $\widehat{f_{\mathfrak{h}}}$ as a tensor product of $\widehat{f_{\mathfrak{h}, \mathfrak{s}_n}} \in \mathcal{S}_c^{\infty}(\mathfrak{s}_n)$ and some function on $\mathfrak{s}_n^{\perp}$. Then we have*

- (i) *The $\widehat{f_{\mathfrak{h}, \mathfrak{s}_n}}$ on $\mathfrak{s}_n$ is supported in $\mathfrak{s}_{n,-}$.*
- (ii) *Let $A = (A_{ij}) \in \mathfrak{s}_n$ be in the support of $\widehat{f_{\mathfrak{h}, \mathfrak{s}_n}}$. Assume further that the absolute values $|\text{tr} \wedge^i A|$ ($i = 1, 2, \dots, n$) are bounded by a constant c . Then the norm of the last column $v_n = (A_{in})_{i=1}^n$ must be bounded (i.e., $\|v_n\|$ is bounded) by some constant depending only on c , n , and the norms of the first $n-1$ columns $\|v_i\|$, $1 \leq i \leq n-1$.*
- (iii) *The intersection of the support of $\widehat{f_{\mathfrak{h}}}$ and the relative nilpotent cone $\mathcal{N} \subset \mathfrak{s}$ is in the H_n -orbit of ξ_- .*

Proof. Suppose that f is (m, m', r) -admissible of the form f^{Ψ} . By the admissibility, there exists some integer $m > 0$ such that

- if A is in the support of $\widehat{f_{\mathfrak{h}}}|_{\mathfrak{s}}$, then $A = (A_{ij})_{i,j=1}^n$ satisfies

$$(8.5) \quad |A_{ij}/A_{i,i-1}| < |\varpi^m|, \quad 1 \leq j < n, j \neq i-1, 1 \leq i \leq n,$$

- $\widehat{f_{\mathfrak{h}}}|_{\mathfrak{s}}$ is invariant under left and right multiplication of $1 + \varpi^m M_{n-1}(\mathcal{O}_F)$ (considered as subgroup of $H_{n-1}(F)$, hence of $H_n(F)$).

- $|A_{i,i-1}|$ are fixed non-zero constants depending only on m .

To show the first part, we may divide the i -th row by $A_{i,i-1}$ and rename the resulted matrix by $\tilde{A}$ which lies in $\mathfrak{h}_n$. It suffices to show $\tilde{A} \in \mathfrak{h}_{n,-}$. By the inequality above, we have

$$\tilde{A} \equiv \begin{pmatrix} * & * & * & * \\ 1 & * & * & * \\ 0 & 1 & * & * \\ 0 & 0 & 1 & * \end{pmatrix} \pmod{\varpi \mathcal{O}_F}.$$

Then it is clear that $\tilde{A} \in \mathfrak{h}_{n,-}$. This completes the proof of the first part.

To show the second part, by the invariance of $\hat{f}_{\mathfrak{h}, \mathfrak{s}_n}$ under $1 + \varpi^m M_{n-1}(\mathcal{O}_F)$, for A in the support of $\hat{f}_{\mathfrak{h}, \mathfrak{s}_n}$, there exists $h \in 1 + \varpi^m M_{n-1}(\mathcal{O}_F)$ such that hAh^{-1} remains in the support and is of the form

$$\begin{pmatrix} * & * & * & * \\ a_1 & * & * & * \\ 0 & a_{n-2} & * & * \\ 0 & 0 & a_{n-1} & * \end{pmatrix}.$$

In particular, the absolute value $|a_i|$'s are non-zero fixed constants. Then we may prove *that the entries of the last column of hAh^{-1} is a polynomial of the entries of the first $n-1$ columns of hAh^{-1} possibly inverting a_i 's and the coefficients of characteristic polynomial of A .*⁷ This shows the last column of hAh^{-1} satisfies the the assertion of (ii). Since h is in a compact group $1 + \varpi^m M_{n-1}(\mathcal{O}_F)$, the assertion for A follows too.

We now prove the third part. Let $X_0 = \begin{pmatrix} A_0 & * \\ * & * \end{pmatrix} \in \mathfrak{s}$ be in the support and H_n -unipotent. We need to show $X_0 \in \mathfrak{s}_{n+1,-}$. The H_n -unipotence implies that $A_0^n = 0$. Since obviously A_0 must also be in the support of $\hat{f}_{\mathfrak{h}, \mathfrak{s}_n}$, by the assertion of part (ii) we may assume the last column has norm bounded by a constant depending only on n, m , the fields E/F . By the condition on the support of $\widehat{\phi_n}$, we have $|X_{n+1,j}/X_{n+1,n}| \leq |\varpi^{m'}|$ for some large m' . When the support of f_{n+1} is small enough (equivalently the integer r is large enough), we clearly have for $h \in 1 + \varpi^m M_{n,n}(\mathcal{O}_F)$:

$$\hat{f}_{\mathfrak{h}}(hX_0h^{-1}) = \hat{f}_{\mathfrak{h}}(X_0).$$

Now we may choose a suitable $h \in 1 + \varpi^m M_{n,n}(\mathcal{O}_F)$ such that the last column of $X_1 = hX_0h^{-1}$ is of the form

$$(0, 0, \dots, 0, a_n, *).$$

Then it suffices to show that $X_1 = \begin{pmatrix} A_1 & * \\ * & * \end{pmatrix} \in \mathfrak{s}_{n+1,-}$. Since X_1 remains in the support of $\hat{f}_{\mathfrak{h}}$, the matrix A_1 is in the support of $\hat{f}_{\mathfrak{h}, \mathfrak{s}_n}$. From the first part of the lemma, it follows that $A_1 \in \mathfrak{s}_{n-}$. This completes the proof.

⁷To see this, let the last column of hAh^{-1} be $(A_{ni})_{i=1,\dots,n}$. Then we may go by backward induction on i : if $i = n$, clearly the first $n-1$ columns and the $\text{tr} A$ can generate A_{nn} ; in general, we may show that A_{ni} is a polynomial of the first $n-1$ columns possibly inverting a_i 's and $\text{tr } \wedge^j A$ for $j = 1, \dots, n-i+1$.

□

We now are ready to prove the local constancy property. This can be viewed as a truncated form of Shalika germ expansion (around zero) of the orbital integral of $\widehat{f}_{\mathfrak{h}}$ for admissible f .

Proposition 8.7. *If f is sufficiently admissible, then there is a small neighborhood $\mathcal{Z}$ of $X = 0$ such that $\eta(\Delta_-(X))O(X, \widehat{f}_{\mathfrak{h}})$ is a constant for regular semisimple X . And the constant is given by the regular unipotent orbital integral:*

$$O(X, \widehat{f}_{\mathfrak{h}}) = \eta(\Delta_-(X/\tau))\mu_{\xi_-}(\widehat{f}_{\mathfrak{h}}).$$

Moreover, for any fixed compact neighborhood $\mathcal{Z}$ of $X = 0$ in $\mathfrak{s}$, there exists an admissible f such that the above equality holds for all regular semisimple X in $\mathcal{Z}$ and the RHS does not vanish.

Proof. We first assume that $\widehat{f}_{\mathfrak{h}}$ is supported in $\mathfrak{s}_{n+1,-}$. Then it follows from Prop. 6.9 (or instead, its transpose) that

$$\eta(\Delta_-(X/\tau))O(X, f) = \mu_{\xi_-}(f)$$

is a constant in a small neighborhood of zero. In general, we may choose a function $f_0 \in \mathcal{C}_c^\infty(\mathfrak{s})$ supported in $\mathfrak{s}_{n+1,-}$ with the same regular unipotent orbital integral as $\widehat{f}_{\mathfrak{h}}$:

$$\mu_{\xi_-}(f_0) = \mu_{\xi_-}(\widehat{f}_{\mathfrak{h}}).$$

By the previous lemma, the functions f_0 and $\widehat{f}_{\mathfrak{h}}$ have the same image in $\mathcal{S}(\mathcal{N})_{H_n, \eta}$. Equivalently, for any (H_n, η) -invariant distribution T supported in $\mathcal{N}$, $T(f_0 - \widehat{f}_{\mathfrak{h}}) = 0$. We claim that $O(X, f_0 - \widehat{f}_{\mathfrak{h}})$ is zero when X is in a small neighborhood of $0 \in \mathfrak{s}$. Indeed, this follows from a variant of Bernstein-Zelvinsky localization principle (cf., [28, p.513, Lemma 27.1]): for $\pi : \mathfrak{s}_{n+1} \rightarrow \mathcal{X}$ we have

$$\lim_U \mathcal{C}_c^\infty(\pi^{-1}(U))_{H_n, \eta} \simeq \mathcal{C}_c^\infty(\pi^{-1}(0))_{H_n, \eta},$$

where U runs over the set of open neighborhoods of 0 in $\mathcal{X}$. Since $\mathcal{N} = \pi^{-1}(0)$ and the image of $(f_0 - \widehat{f}_{\mathfrak{h}})$ in $\mathcal{C}_c^\infty(\pi^{-1}(0))_{H_n, \eta}$ is zero, there exists some neighborhood U of 0 such that the image of $(f_0 - \widehat{f}_{\mathfrak{h}})$ in $\mathcal{C}_c^\infty(\pi^{-1}(U))_{H_n, \eta}$ is zero. This completes the proof of the first part.

To show the second part, we start with any (m_0, m'_0, r_0) -admissible

$$f' = f^\Psi = f_n^{\varphi_{n-1}, \phi_{n-1}} \otimes f_{n+1}^{\phi_n^\vee}$$

with the following property:

- (1) The regular unipotent orbital integral $\mu_{\xi_-}(\widehat{f}'_{\mathfrak{h}}) \neq 0$.

(2) The function $\phi_n \in \mathcal{C}(M_{1,n}(E))_{m'_0}^\bullet$ is chosen such that

$$\widehat{\phi}_n(x_1, \dots, x_n) = \widehat{\phi}_n^n(x_n) 1_{\varpi_E^{m'_0} M_{1,n-1}(\mathcal{O}_E)}(x_1/x_n, \dots, x_{n-1}/x_n),$$

where $\phi_n^n \in \mathcal{C}_c^\infty(E)$ is a suitable function which we will keep fixed.

We then assume that the local constancy holds for a non-empty open neighborhood $\mathcal{Z}_0$ of 0. We claim that *if we increase m'_0 sufficiently and then let r'_0 be sufficiently large, we can maintain the local constancy on $\mathcal{Z}_0$* . Indeed, since the orbital integral depends only on H_n -orbits, it suffices to consider orbital integral of regular semisimple elements of the form ${}^t\sigma(a, b)$, the transpose of $\sigma(a, b)$ (cf. §6 for definition) for (a, b) in the image of $\mathcal{Z}_0$ in $\mathcal{X}$. We may replace the integral over $H_n(F)$ by the integral over an open dense subset

$$h = ku, \quad u = \begin{pmatrix} 1_{n-1} & \\ v_{n-1} & 1 \end{pmatrix}, v_{n-1} \in M_{1,n-1}(F)$$

and k is in P_n consisting of $\begin{pmatrix} * & * \\ 0_{n-1} & * \end{pmatrix} \in H_n$. By definition we have

$$O(X, \widehat{f}'_n) = \int_{M_{1,n-1}(F)} \int_{P_n} \widehat{f}'_n(u^{-1}k^{-1}Xku) \eta(h) dk du$$

where dk is the left invariant Haar measure on P_n . For $X = {}^t\sigma(a, b)$, the last row of $h^{-1}Xh$ is of the form $a_n(v_{n-1}, 1)$ where $a_n \in F^\times$ is the last entry of k . By the assumption on ϕ_n (or rather, $\widehat{\phi}_n$), we see that the integrand is non-zero only if $\|v_{n-1}\| \leq |\varpi|^{m'_0}$. Moreover, our $X = {}^t\sigma(a, b)$ lies in a bounded domain $\mathcal{Z}_0$. If $\widehat{f}'_n(u^{-1}k^{-1}Xku) \neq 0$, then $u^{-1}k^{-1}Xku$ lies in the support of $\widehat{f}'_n$. By part (ii) of Lemma 8.6, the last row of $u^{-1}k^{-1}Xku$ must be in a bounded domain with radius depending only on m_0 , but not on m'_0 nor r_0 . Write $X = \begin{pmatrix} A & * \\ * & * \end{pmatrix}$, where $A \in M_n(F)$. It follows that when m'_0 becomes sufficiently large, for $X = {}^t\sigma(a, b) \in \mathcal{Z}_0$, we have $\widehat{f}'_n(u^{-1}k^{-1}Xku) = \widehat{f}'_n(k^{-1}Ak)$ (also cf. the proof of Lemma 8.8). We thus obtain

$$O(X, \widehat{f}'_n) = \left(\int_{M_{1,n-1}(F)} 1_{\varpi_E^{m'_0} M_{1,n-1}(\mathcal{O}_E^-)}(v_{n-1}) dv_{n-1} \right) \int_{P_n} \widehat{f}'_n(k^{-1}Xk) \eta(k) dk.$$

As the first factor is a constant and the second factor does not see m'_0 , the claim now follows.

We return to the proof of the second part. By the compactness of $\mathcal{Z}$, we may assume that $t\mathcal{Z} \subset \mathcal{Z}_0$ for some $t \in F^\times$ with sufficiently small norm $|t|$. Now we may replace ϕ_n by $\phi_{n,t}$ which is of the same form but we change m'_0 to a sufficiently large m'_1 and change

$$\phi_{n,t}^n(x_n) = \phi_{n,t}^n(x_n/t).$$

We also choose $\varphi_{n-1,t}, \phi_{n-1,t}$ such that $\Psi_t = (\varphi_{n-1}, \phi_{n-1}, \phi_n)$ has the property

$$\Psi_t^-(p) = \Psi^-(p/t).$$

Note that $(\varphi_{n-1}, \phi_{n-1})$ remains m_0 -admissible (but with Λ replaced by another lattice $t\Lambda$, cf. §7 for definition). Now we let $m'_1 > m_1$ be sufficiently large to fulfill the second condition in the definition of admissible function (cf. §8, Definition 8.1). Then with sufficiently large r_1 such that $f'_i = 1 + \varpi^{r_1} M_i(\mathcal{O}_E)$, $i = n, n+1$, the function

$$f'_t = f_n^{\varphi_{n-1,t}, \phi_{n-1,t}} \otimes f_{n+1}^{\phi_n^\vee, t}$$

is (m_0, m'_1, r_1) -admissible (with smaller support than f'). Then we claim that with the new choice, the local constancy holds on $\mathcal{Z}_0/t$ which contains $\mathcal{Z}$. Indeed, we may choose r_0, r_1 sufficiently large such that $r_1 = r_0 + v(t)$. Then we see that

$$f'_{t, \mathfrak{q}}(X) = f'_\mathfrak{q}(X/t).$$

Then their Fourier transforms satisfy

$$\widehat{f'_{t, \mathfrak{q}}}(X) = c_t \widehat{f'_\mathfrak{q}}(Xt).$$

for some constant $c_t \neq 0$ depending only on t . Then clearly

$$O(X, \widehat{f'_{t, \mathfrak{q}}}) = c_t O(tX, \widehat{f'_\mathfrak{q}}).$$

The local constancy of $O(X, \widehat{f'_\mathfrak{q}})$ holds on $\mathcal{Z}_0$ as shown earlier. This completes the proof of the second part of the proposition. $\square$

A formula for the regular unipotent orbital integral. We now consider the regular unipotent orbital integral of Fourier transform $\widehat{f'_\mathfrak{q}}^\Psi$ of $f'_\mathfrak{q}^\Psi$ on $\mathfrak{s}$. Recall that the regular unipotent orbit is given by

$$\xi_- = \xi_{n+1, -} = \tau \begin{pmatrix} 0 & 0 & \dots & 0 \\ 1 & 0 & 0 & \dots \\ \dots & 1 & 0 & 0 \\ 0 & \dots & 1 & 0 \end{pmatrix} \in \mathfrak{s}.$$

Lemma 8.8. *Fix Π and assume that $f^\Psi = f_n^{\varphi_{n-1}, \phi_{n-1}} \otimes f_{n+1}^{\phi_n^\vee}$ is admissible for Π . Then we have*

$$\mu_{\xi_-}(\widehat{f'_\mathfrak{q}}^\Psi) = c(\Psi^+) |\overline{\delta}_{n,E}(\epsilon_n)|^{-1/2} \prod_{i=1}^{n-1} \phi_i^{\prime -}(0) \int_{A_n(F) N_{n,-}(F)} \prod_{i=1}^n \widehat{\phi}_i^-(-a_i(v_{i-1}, 1)\tau) |\overline{\delta}_n(a)|^{-1} \eta(a) d^*adv,$$

where $v_i \in M_{i,1}(F)$ and $\overline{\delta}_n(a)$ is the same as before.

Proof. Without loss of generality we may assume $c(\Psi^+) = 1$. Assume that f^Ψ is (m, m', r) -admissible. We denote $f' = (f_{\mathfrak{h}}^\Psi)^\vee$. By definition we have

$$\mu_{\xi_-}(\widehat{f_{\mathfrak{h}}^\Psi}) = \int_{H_n(F)} \widehat{f_{\mathfrak{h}}^\Psi}(h^{-1}\xi_{n+1,-}h)\eta(h)dh = \int_{H_n(F)} f'(-h^{-1}\xi_{n+1,-}h)\eta(h)dh.$$

When the support of f_{n+1} is small enough, by the previous Lemma 8.5 the function f' on $\mathfrak{s}$ is of the form $\Psi^- \otimes \phi_n'^- \otimes \phi_{n+1}^-$ corresponding to decomposition

$$\mathfrak{s} = \mathfrak{p}^- \oplus M_{1,n}(E^-) \oplus M_{1,n+1}(E^-)$$

where $\phi_i'^-$ is a multiple of $1_{M_{1,i}(\mathcal{O}_E^-)}$ normalized so that $\int \phi_i'^-(z)dz = 1$ or equivalently $\widehat{\phi_i'^-}(0) = 1$, for $i = n, n+1$. The Fourier transform takes the form $\widehat{\Psi}^- \otimes \widehat{\phi_n'^-} \otimes \widehat{\phi_{n+1}^-}$ where $\widehat{\Psi}^-$ is a function on the transpose of $\mathfrak{p}^-$. The way we choose the notation is such that the function $f_{\mathfrak{h}}^\Psi$ is of the following form:

$\phi_1'^-$	ϕ_1^-	ϕ_2^-	$\cdots$	ϕ_n^-
$\phi_2'^-$				
$\vdots$				
$\vdots$				
$\phi_{n+1}'^-$				

We also write $f_{\mathfrak{h}}^\Psi$ as the tensor product

$$\varphi_n^- \otimes \phi_n^- \otimes \phi_{n+1}'^-,$$

where $\varphi_n^- \in \mathcal{C}_c^\infty(\mathfrak{s}_n)$. Therefore we may consider the Fourier transform $\widehat{f_{\mathfrak{h}}^\Psi}$ as $\widehat{\varphi_n^-} \otimes \widehat{\phi_n^-} \otimes \widehat{\phi_{n+1}'^-}$, or as the tensor product of $\bigotimes_{i=1}^{n+1} \widehat{\phi_i'^-}$ and $\bigotimes_{i=1}^n \widehat{\phi_i^-}$. Most importantly for our computation, *the function $\widehat{\varphi_n^-}$ on $\mathfrak{s}$ is invariant under left and right multiplication by $1 + \varpi^m M_n(\mathcal{O}_F)$.* Back to the orbital integral, we may write $h \in H'_n(F)$ as

$$h = a_n k v, \quad k \in N_{n-1} H_{n-1}, v = \begin{pmatrix} 1_{n-1} & \\ & 1 \end{pmatrix}, a_n \in F^\times.$$

Then the last row of $h^{-1}\xi_-h$ is of the form

$$a_n(v_{n-1}, 1)\tau.$$

Then by the condition on the support of f' (more precisely, of $\widehat{\phi_n^-}$ or $\widehat{\phi_n'^-}$ since $\phi_n \in \mathcal{C}_c^\infty(M_{n,1}(E))_{m'}^\dagger$), we may assume that $\|v_{n-1}\| \leq |\varpi^{m'}|$ is small. By the part (ii) of Lemma 8.6, if m' is large enough we have

$$f'(-h^{-1}\xi_{n+1,-}h) = \widehat{\varphi_n^-}(-k^{-1}\xi_{n,-}k)\widehat{\phi_n^-}(-a_n(v_{n-1}, 1)\tau)\widehat{\phi_{n+1}'^-}(0)$$

if the LHS is not zero (i.e., at least $h^{-1}\xi_{n+1,-}h$ lies in the support.) Note that $\widehat{\phi'_{n+1,-}}(0) = 1$ by our normalization. We now may repeat the process and utilize the invariance of $\widehat{\varphi_n^-}$ under $1 + \varpi^m M_{n,n}(\mathcal{O}_F)$. To state the outcome we let

$$h = uav, a = \begin{pmatrix} a_1 a_2 \dots a_n & & \\ & \ddots & \\ & & a_n \end{pmatrix}, u \in N_n(F),$$

and write $v \in N_{n-}(F)$ as the product of $\begin{pmatrix} 1_{n-i} & \\ v_{n-i} & 1 \end{pmatrix} \in N_{n-i+1,-}(F) \subset N_{n-}(F)$ for $1 \leq i \leq n-1$. Note that $u^{-1}\xi_{n-}u$ is of the form

$$\tau \begin{pmatrix} * & * & * & * \\ 1 & * & * & * \\ 0 & 1 & * & * \\ 0 & 0 & 1 & * \end{pmatrix}.$$

If $v \in N_{n-i-1,-}(F)$, then the conjugation $v^{-1}u^{-1}\xi_{n-}uv$ does not change the last i rows below the diagonal of $u^{-1}\xi_{n-}u$. We now view $\bigotimes_{i=1}^n \widehat{\phi'_i}$ as a function on the set of upper triangular elements and then consider it as a function on $\mathfrak{s}$ via the natural projection—denoted by γ —from $\mathfrak{s}$ to the upper triangular elements. Then the integral $\mu_{\mathcal{O}_0}(\widehat{f_{\mathfrak{h}}^\Psi})$ is equal to

$$\int \left(\bigotimes_{i=1}^{n+1} \widehat{\phi'_{n-i}} \right) (-(ua)^{-1}\xi_-ua) du \prod_{i=1}^n \widehat{\phi'_i}(-a_i(v_{i-1}, 1)\tau) |\delta_n(a)|^{-1} \eta(a) d^*a dv,$$

where $u \in N_n(F), v \in N_{n-}(F)$. When r is large enough, the function $\widehat{\phi'_n}$ is a constant in a very large neighborhood of 0. By Lemma 6.2, the last column of $u^{-1}\xi_{n-}u$ is a polynomial of the first $n-1$ columns. By the same lemma, after a suitable substitution we may write the integral over u

$$\int \left(\bigotimes_{i=1}^n \widehat{\phi'_i} \right) \circ \gamma((ua)^{-1}\xi_-ua) du = \int \left(\bigotimes_{i=1}^n \widehat{\phi'_i} \right) \circ \gamma(a^{-1}u'a) du',$$

where the integral of u' is over the space of elements of the form

$$\tau \begin{pmatrix} * & * & * & 0 \\ 1 & * & * & 0 \\ 0 & 1 & * & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix},$$

and the measure du' is induced by du . We now make the substitution $u' \mapsto au'a^{-1} = Ad(a)u'$ (equivalently, $u'_{ij} \mapsto u'_{ij} \prod_{i < \ell \leq j} a_\ell$ for $1 \leq i \leq n-1$): This yields

$$\begin{aligned} & \int \left(\bigotimes_{i=1}^n \widehat{\phi_i'^-} \right) \circ \gamma(a^{-1}u'a) du' \\ &= \det(Ad(a) : \mathfrak{n}_{n-1}) \int \left(\bigotimes_{i=1}^n \widehat{\phi_i'^-} \right) \circ \gamma(u') du' \\ &= \det(Ad(a) : \mathfrak{n}_{n-1}) |\tau|_E^{-(1+2+\dots+(n-2))/2} \prod_{i=1}^n \phi_i'^-(0). \end{aligned}$$

(Or more explicitly $\det(Ad(a) : \mathfrak{n}_{n-1}) = \prod_{i=1}^{n-1} |a_i|^{(i-1)(n-i)}$.) Here the factor

$$|\tau|_E^{(1+2+\dots+(n-2))/2} = |\bar{\delta}_{n,E}(\epsilon_n)|^{1/2}$$

comes from the difference between the measure on $F\tau$ and E^- . Return to $\mu_{\mathcal{O}_0}(\widehat{f}_{\mathfrak{h}})$:

$$|\bar{\delta}_{n,E}(\epsilon_n)|^{1/2} \prod_{i=1}^n \phi_i'^-(0) \int_{A_n(F)N_{n,-}(F)} \prod_{i=1}^n \widehat{\phi_i'^-}(-a_i(v_{i-1}, 1)\tau) |\bar{\delta}_n(a)|^{-1} \eta(a) d^*adv.$$

(Or more explicitly, $\bar{\delta}_n(a) = \det(Ad(a) : \mathfrak{n}_n) / \det(Ad(a) : \mathfrak{n}_{n-1}) = \prod_{i=1}^{n-1} |a_i|^{n-i}$.) We finally note that $\phi_n'^-(0) = 1$ to complete the proof. $\square$

Proof of Theorem 8.4. This follows from Prop. 8.3 and Lemma 8.8. We decompose $\varphi_{n-1}, \phi_{n-1}$ as in Lemma 7.1 and similarly ϕ_n . Note that

$$\begin{aligned} & \int \prod_{i=0}^{n-1} \widehat{\phi_{n-i}^-}(-y_i(z_{n-i-1}, 1)\tau) |\bar{\delta}_n(y')|^{-1} \eta(y') d^*y' dz, \\ &= \prod_{i=0}^{n-1} \widehat{\phi_{n-i}^+}(0) \int \prod_{i=0}^{n-1} \widehat{\phi_{n-i}^-}(-y_i(z_{n-i-1}, 1)\tau) |\bar{\delta}_n(y')|^{-1} \eta(y') d^*y' dz \\ &= \prod_{i=1}^n \int \phi_i^+(x_i) dx_i \int \prod_{i=0}^{n-1} \widehat{\phi_{n-i}^-}(-y_i(z_{n-i-1}, 1)\tau) |\bar{\delta}_n(y')|^{-1} \eta(y') d^*y' dz. \end{aligned}$$

and

$$\int_{B_{n-1,-}(F)} \phi'(b) db = \prod_{i=1}^{n-1} \phi_i'^-(0) \prod_{i=1}^{n-1} \int \phi_i'^+(b_i) db_i.$$

Now it is clear that

$$c(\Psi^+) = \prod_{i=1}^n \int \phi_i^+(x_i) dx_i \prod_{i=1}^{n-1} \int \phi_i'^+(b_i) db_i.$$

Note that

$$|\tau|_E^{(d_n+d_{n+1})/2} = |\delta_{n-1,E}(\epsilon_{n-1})|^{1/2} |\delta_{n,E}(\epsilon_n)|^{1/2} = |\delta_{n-1,E}(\epsilon_{n-1})| |\bar{\delta}_{n,E}(\epsilon_n)|^{1/2}.$$

Then Theorem 8.4 follows by comparison with Lemma 8.8.

9 Local character expansion in the unitary case

Three ingredients from [50]. Let F be a non-archimedean local field of characteristic zero. We need to recall the main local results of [50]. There are two isomorphism classes of Hermitian space W_1, W_2 of dimension n . Denote by $H_{W_i} = U(W_i)$ the unitary group. We let $V_i = W_i \perp Ee$ be the orthogonal sum of W_i and a one-dimensional space Ee with norm $\langle e, e \rangle = 1$. Denote by $\mathfrak{u}_i$ the Lie algebra of $U(V_i)$. We have a bijection of regular semisimple orbits

$$H_n(F) \backslash \mathfrak{s}(F)_{rs} \simeq H_{W_1}(F) \backslash \mathfrak{u}_1(F)_{rs} \coprod H_{W_2}(F) \backslash \mathfrak{u}_2(F)_{rs}$$

A regular semisimple $X \in \mathfrak{s}$ matches some $Y \in \mathfrak{u}_i$ if and only if

$$\eta(\Delta(X/\tau)) = \eta(\text{disc}(W_i)),$$

where $\text{disc}(W_i) \in F^\times / NE^\times$ is the discriminant of W_i . For $f' \in \mathcal{C}_c^\infty(\mathfrak{s})$ and the pair (f_1, f_2) , $f_i \in \mathcal{C}_c^\infty(\mathfrak{u}_i)$, we say that f' match (f_1, f_2) , if for all matching regular semisimple $X \in \mathfrak{s}, Y \in \mathfrak{u}_i$, we have

$$\eta'(\Delta_+(X))O(X, f') = O(Y, f_i).$$

Here η' is a fixed choice of character $E^\times \rightarrow \mathbb{C}^\times$ with restriction $\eta'|_{F^\times} = \eta$. We also say $f_1 \in \mathcal{C}_c^\infty(\mathfrak{u}_1)$ and $f' \in \mathcal{C}_c^\infty(\mathfrak{s})$ match if there exists $f_2 \in \mathcal{C}_c^\infty(\mathfrak{u}_2)$ such that the pair (f_1, f_2) and f' match.

Analogous to the general linear case, to a function in a small neighborhood of $1 \in G = U(W) \times U(V)$, we would like to associate a function on the Lie algebra $\mathfrak{u}$ of $U(V)$. For $f = f_n \otimes f_{n+1} \in \mathcal{C}_c^\infty(G)$, we let $\tilde{f}$ be the function on $U(V)$ defined by

$$\tilde{f}(g) := \int_{U(W)} f_n(h) f_{n+1}(hg) dh, \quad g \in U(V).$$

If f is supported in a small neighborhood Ω of 1 in G , then $\tilde{f}$ is supported in a small neighborhood $\tilde{\Omega}$ of 1 in $U(V)$. Since the Cayley map $\mathfrak{c} : \mathfrak{u} \rightarrow U(V)$ is a local homeomorphism around $0 \in \mathfrak{u}$, we may denote $\omega = \mathfrak{c}^{-1}(\tilde{\Omega}) \simeq \tilde{\Omega}$ and to $\tilde{f}$ we associate a function denoted by $f_{\mathfrak{h}} = \mathfrak{c}^{-1}(\tilde{f})$ on $\mathfrak{u}$ supported in ω . To connect the smooth transfer on the groups to the one on Lie algebras, we need:

Lemma 9.1. *Let $f' \in \mathcal{C}_c^\infty(G'(F))$ and $f_i \in \mathcal{C}_c^\infty(U(W_i) \times U(V_i))$ be matching functions (in the sense of §4) with small support around the identity. Then the functions $\eta(2^n) f'_{\mathfrak{h}} \in \mathcal{C}_c^\infty(\mathfrak{s})$ and $f_{i,\mathfrak{h}} \in \mathcal{C}_c^\infty(\mathfrak{u}_i)$ match.*

Proof. The support condition insures that the associated functions $f'_{\mathfrak{h}}, f_{i,\mathfrak{h}}$ are well-defined. Then it remains to show the transfer factor are compatible:

$$\eta'(\Delta_+(X))O(X, f'_{\mathfrak{h}}) = \Omega(g)O(g, f'),$$

where $\nu(g) = \mathfrak{c}(X) \in S_{n+1}(F)$. This follows from the proof of [50, Lemma 3.5]. $\square$

In [50, Theorem 2.6] the following result is proved:

Theorem 9.2. *For any $f \in \mathcal{C}_c^\infty(\mathfrak{u})$ there exists a matching $f' \in \mathcal{C}_c^\infty(\mathfrak{s})$ and conversely.*

Moreover we have [50, Theorem 4.17]:

Theorem 9.3. *If f and f' match, then so do $\epsilon(1/2, \eta, \psi)^{n(1+n)/2} \hat{f}$ and $\hat{f}'$.*

An important ingredient of the proof of both theorems above is a local trace formula on Lie algebra ([50, Theorem 4.6]). Now we only need the one in the unitary case.

Theorem 9.4. *For $f_1, f_2 \in \mathcal{C}_c^\infty(\mathfrak{u})$, we have*

$$\int_{\mathfrak{u}} f_1(X)O(X, \hat{f}_2)dX = \int_{\mathfrak{u}} O(X, \hat{f}_1)f_2(X)dX,$$

where the integrals are absolutely convergent.

A hypothesis. We now return to the spherical character in the unitary case. Let π be an irreducible admissible representation of $G = U(W) \times U(V)$. We use the measure on $U(V)$ determined by the self-dual measure on $\mathfrak{u}$ via the Cayley map. Consider

Hypothesis ($\star$): there exists a small neighborhood $\Omega \subset G$ of 1 and a function $\Phi \in \mathcal{C}_c^\infty(\mathfrak{u})$ such that

$$(9.1) \quad \Phi(0) = 1,$$

and for all $f \in \mathcal{C}_c^\infty(\Omega)$,

$$J_\pi(f) = \int_{\mathfrak{u}} f_{\mathfrak{h}}(X)O(X, \Phi)dX.$$

Theorem 9.5. *Assume that π is tempered and $\text{Hom}_H(\pi, \mathbb{C}) \neq 0$. Let Φ be a matrix coefficient of π such that*

$$\int_H \Phi(h)dh = 1.$$

1. *The distribution J_π is representable by the orbital integral of Φ*

$$O(x, \Phi).$$

Moreover, it is a bi- H -invariant function that is locally L^1 on G .

2. Assume further that one of the following two is satisfied

- The representation π is supercuspidal.
- The group H is compact.

Then the function $x \mapsto O(x, \Phi)$ is locally constant on the regular semisimple locus.

Proof. When π is tempered and $\text{Hom}_H(\pi, \mathbb{C}) \neq 0$, there exists Φ such that $\int_H \Phi(h)dh \neq 0$. Up to a scalar multiplication, we may assume that $\int_H \Phi(h)dh = 1$. Then the first part is proved in [24]. For the second part, it is clear if H is compact. If it is supercuspidal, Φ has compact support and it follows from a general fact proved in [50] that the orbital integral is locally constant on the regular semisimple locus. $\square$

Proposition 9.6. *Assume that $\text{Hom}_H(\pi, \mathbb{C}) \neq 0$. If the group $H = U(W)$ is compact or π is supercuspidal, then π verifies Hypothesis $(\star)$.*

Proof. Assume that π is supercuspidal. For a small neighborhood of $0 \in \mathfrak{u}$ we will identify it with a neighborhood of $1 \in U(V)$ via the Cayley map. We choose an open and closed H -invariant neighborhood that contains the above chosen neighborhood of $1 \in U(V)$. To truncate Φ , we restrict the function Φ to this neighborhood and denote the corresponding function on $\mathfrak{u}$ still by Φ . It is important to note that we still have

$$(9.2) \quad \Phi(0) = 1.$$

Moreover the measure on $\mathfrak{u}$ is transferred to the measure on $U(V)$. Then $\Phi \in \mathcal{C}_c^\infty(\mathfrak{u})$ and for all function $f \in \mathcal{C}_c^\infty(G)$ with small support around 1,

$$J_\pi(f) = \int_{\mathfrak{g}} f_{\mathfrak{h}}(X) O(X, \Phi) dX.$$

If H is compact (so that $\dim W \leq 2$ or $E/F = \mathbb{C}/\mathbb{R}$), then there is a non-zero vector $\phi_0 \in \pi$ fixed by H . Then for all f , we have

$$J_\pi(f) = \text{vol}(H) \int_G f(g) \langle \pi(g)\phi_0, \phi_0 \rangle dg$$

for a norm one $\phi_0 \in \pi^H$. Set $\Phi(g) = \text{vol}(H)^{-1} \langle \pi(g)\phi_0, \phi_0 \rangle$. Then the same truncation as above completes the proof. $\square$

We say that f is admissible if there is an admissible f' matching f .

Theorem 9.7. *Assume that π verifies Hypothesis $(\star)$. there exists admissible functions $f \in \mathcal{C}_c^\infty(G(F))$ and matching function $f' \in \mathcal{C}_c^\infty(G'(F))$ such that*

$$J_\pi(f) = (\eta'(\tau)/\epsilon(1/2, \eta, \psi))^{n(n+1)/2} \eta(\text{disc}(W)) \widehat{\mu_{\xi_-}}(f'_1) \neq 0.$$

Proof. First we apply the local trace formula (Theorem 9.4)

$$\int_{\mathfrak{u}} f_{\mathfrak{q}}(Y) O(Y, \Phi) dY = \int_{\mathfrak{u}} O(Y, \hat{f}_{\mathfrak{q}}) \check{\Phi}(Y) dY,$$

where $\check{\Phi}$ is the inverse of the Fourier transform. Let f' be an admissible function matching f . Then by the compatibility between Fourier transform and smooth transfer (Theorem 9.3)

$$\epsilon(1/2, \eta, \psi)^{n(n+1)/2} O(Y, \hat{f}_{\mathfrak{q}}) = \eta'(\Delta_+(X)) O(X, \hat{f}'_{\mathfrak{q}})$$

for matching regular semisimple X and Y . Since $\check{\Phi}$ has compact support, by Prop. 8.7, we may choose a sufficiently admissible functions f' such that

$$O(Y, \hat{f}_{\mathfrak{q}}) = (\eta'(\tau)/\epsilon(1/2, \eta, \psi))^{n(n+1)/2} \eta(\Delta(X/\tau)) \mu_{\xi_-}(\hat{f}'_{\mathfrak{q}}) \neq 0$$

is a constant for all regular semisimple Y in the support of $\check{\Phi}$. We know that for regular semisimple X , its transpose X^t is in the same H_n -orbit and $X^t = hXh^{-1}$ for a unique $h \in H_n(F)$ with

$$\eta(h) = \eta(\Delta_+(X)/\Delta_-(X)) = \eta(\Delta(X/\tau)) = \eta(\text{disc}(W)).$$

Therefore for all regular semisimple Y in the support of $\check{\Phi}$ we have

$$O(Y, \hat{f}_{\mathfrak{q}}) = (\eta'(\tau)/\epsilon(1/2, \eta, \psi))^{n(n+1)/2} \eta(\text{disc}(W)) \mu_{\xi_-}(\hat{f}'_{\mathfrak{q}}) \neq 0.$$

We obtain

$$\begin{aligned} J_{\pi}(f) &= \int_{\mathfrak{u}} O(X, \hat{f}_{\mathfrak{q}}) \check{\Phi}(X) dX \\ &= (\eta'(\tau)/\epsilon(1/2, \eta, \psi))^{n(n+1)/2} \eta(\text{disc}(W)) \mu_{\xi_-}(\hat{f}'_{\mathfrak{q}}) \cdot \int_{\mathfrak{u}} \check{\Phi}(X) dX \\ &= (\eta'(\tau)/\epsilon(1/2, \eta, \psi))^{n(n+1)/2} \eta(\text{disc}(W)) \mu_{\xi_-}(\hat{f}'_{\mathfrak{q}}) \cdot \Phi(0). \end{aligned}$$

By Hypothesis $(\star)$ we have

$$\Phi(0) = 1.$$

The theorem now follows. □

Completion of the proof of Theorem 4.5. It remains to prove the case when v is non-split. Since the ratio $I_{\Pi_v}(f'_v)/J_{\pi_v}(f_v)$ is independent of the choice of f_v , we may choose a suitable admissible function. Then the result follows from Theorem 8.4, Prop. 9.6 and Theorem 9.7.

Concluding remarks Note that we only deal with π_v which appears as a local component of a global π . But we expect Conjecture 4.3 holds in general (as long as Π_v is generic in order to define I_{Π_v}).

We conclude with

Conjecture 9.8. *The spherical character I_{Π} and J_{π} are representable by a locally L^1 function and smooth (local constant in the non-archimedean case) on an open subset.*

There should be a more complete analogue of the Harish–Chandra local character expansion in our relative setting. Moreover, it seems that the spherical characters (if non-zero) should contains as much information as the usual characters of the representation.

References

- [1] Aizenbud, Gourevitch, Rallis, Schiffmann, *Multiplicity one theorems*, Ann. of Math. (2) 172 (2010), no. 2, 1407–1434.
- [2] R. Beuzart-Plessis, *La conjecture locale de Gross-Prasad pour les représentations tempres des groupes unitaires*, arXiv:1205.2987
- [3] Jacquet, Hervé; Chen, Nan. *Positivity of quadratic base change L -functions*. Bull. Soc. Math. France 129 (2001), no. 1, 3390.
- [4] Flicker, Yuval Z. *Twisted tensors and Euler products*. Bull. Soc. Math. France 116 (1988), no. 3, 295–313.
- [5] Flicker, Yuval Z. *On distinguished representations*. J. Reine Angew. Math. 418 (1991), 139–172.
- [6] Wee Teck Gan, Benedict H. Gross and Dipendra Prasad. *Symplectic local root numbers, central critical L -values, and restriction problems in the representation theory of classical groups* Preprint, <http://www.math.harvard.edu/~gross/preprints/>
- [7] Wee Teck Gan, Atsushi Ichino. *On endoscopy and the refined Gross-Prasad conjecture for $(\mathrm{SO}_5, \mathrm{SO}_4)$* . to appear in J. Math Institute Jussieu.
- [8] P. Garrett, *Decomposition of Eisenstein series: Rankin triple products*. Ann. of Math. (2) 125 (1987), no. 2, 209–235.
- [9] Gelbart, Stephen; Jacquet, Herv; Rogawski, Jonathan *Generic representations for the unitary group in three variables*. Israel J. Math. 126 (2001), 173–237.
- [10] Ginzburg, David; Jiang, Dihua; Rallis, Stephen *On the nonvanishing of the central value of the Rankin-Selberg L -functions*. J. Amer. Math. Soc. 17 (2004), no. 3, 679–722

- [11] Ginzburg, David; Jiang, Dihua; Rallis, Stephen *On the nonvanishing of the central value of the Rankin-Selberg L-functions. II.* Automorphic representations, L-functions and applications: progress and prospects, 157–191
- [12] Gross, Benedict H. *Heights and the special values of L-series.* Number theory (Montreal, Que., 1985), 115–187, CMS Conf. Proc., 7, Amer. Math. Soc., Providence, RI, 1987.
- [13] Gross, Benedict H. *On the motive of a reductive group.* Invent. Math. 130 (1997), no. 2, 287–313.
- [14] Gross, Benedict H.; Prasad, Dipendra *On the decomposition of a representation of SO_n when restricted to SO_{n-1} .* Canad. J. Math. 44 (1992), no. 5, 974–1002
- [15] Gross, Benedict H.; Prasad, Dipendra *On irreducible representations of $\mathrm{SO}_{2n+1} \times \mathrm{SO}_{2m}$.* Canad. J. Math. 46 (1994), no. 5, 930–950.
- [16] B. Gross; D. Zagier: *Heegner points and derivatives of L-series.* Invent. Math. 84 (1986), no. 2, 225–320.
- [17] Harish-Chandra, *Harmonic analysis on reductive p-adic groups.* Notes by G. van Dijk. Lecture Notes in Mathematics, Vol. 162. Springer-Verlag, Berlin-New York, 1970
- [18] Harish-Chandra, *Admissible invariant distributions on reductive p-adic groups.* Lie theories and their applications (Proc. Ann. Sem. Canad. Math. Congr., Queen’s Univ., Kingston, Ont., 1977), pp. 281–347. Queen’s Papers in Pure Appl. Math., No. 48, Queen’s Univ., Kingston, Ont., 1978
- [19] M. Harris and S. Kudla, *On a conjecture of Jacquet.* Contributions to automorphic forms, geometry, and number theory, 355–371, Johns Hopkins Univ. Press, Baltimore, MD, 2004.
- [20] M. Harris. *L-functions and periods of adjoint motives*, to appear in Algebra and Number Theory.
- [21] Neal Harris, *A refined Gross–Prasad conjecture for unitary groups*, PhD Thesis, 2011, UCSD.
- [22] Ichino Atsushi. *Trilinear forms and the central values of triple product L-functions.* Duke Math. J. Volume 145, Number 2 (2008), 281–307.
- [23] A. Ichino and T. Ikeda. *On the periods of automorphic forms on special orthogonal groups and the Gross-Prasad conjecture*, Geom. Funct. Anal. 19 (2010), no. 5, 1378–1425.
- [24] A. Ichino and W. Zhang. *Spherical characters for a strongly tempered pair* Appendix to [50].
- [25] Jacquet, H.; Piatetskii-Shapiro, I. I.; Shalika, J. A. *Rankin-Selberg convolutions.* Amer. J. Math. 105 (1983), no. 2, 367–464.

- [26] Hervé Jacquet, S. Rallis. *On the Gross-Prasad conjecture for unitary groups*. in *On certain L-functions*, 205264, Clay Math. Proc., 13, Amer. Math. Soc., Providence, RI, 2011.
- [27] Jacquet, H.; Shalika, J. A. *On Euler products and the classification of automorphic representations. I*. Amer. J. Math. 103 (1981), no. 3, 499–558.
- [28] Kottwitz, Robert E. *Harmonic analysis on reductive p -adic groups and Lie algebras*. Harmonic analysis, the trace formula, and Shimura varieties, 393522, Clay Math. Proc., 4, Amer. Math. Soc., Providence, RI, 2005.
- [29] E. Lapid, S. Rallis *On the non-negativity of $L(1/2, \pi)$ for $SO(2n + 1)$* , Annals of Math. 157 (2003), 891-917.
- [30] E. Lapid *On the non-negativity of Rankin-Selberg L -functions at the center of symmetry*, International Mathematics Research Notices 2003 No.2 65-75.
- [31] C-P. Mok, *Endoscopic classification of representations of quasi-split unitary groups I*, arXiv:1206.0882v1
- [32] Rader, Cary; Rallis, Steve. *Spherical characters on p -adic symmetric spaces*. Amer. J. Math. 118 (1996), no. 1, 91–178.
- [33] Steve Rallis, Grard Schiffmann. *Multiplicity one Conjectures*. arXiv:0705.2168.
- [34] Y. Sakellaridis, A. Venkatesh. *Periods and harmonic analysis on spherical varieties*, arXiv:1203.0039.
- [35] F. Shahidi. *On certain L -functions*. Amer. J. Math., 103(2):297-355, 1981.
- [36] Sun B., Zhu C. *Multiplicity one theorems: the Archimedean case*, Annals Math., to appear.
- [37] Ye Tian, *Congruent numbers and Heegner points*, preprint
- [38] J. Waldspurger: *Sur les valeurs de certaines fonctions L automorphes en leur centre de symétrie*. Compositio Math. 54 (1985), no. 2, 173–242.
- [39] J. Waldspurger: *Une formule integrale relie la conjecture locale de Gross-Prasad*. arXiv:0902.1875.
- [40] J. Waldspurger: *Une formule integrale relie la conjecture locale de Gross-Prasad, 2me partie: extension aux representations temporelles*. arXiv:0904.0314.
- [41] P.-J., White. *Tempered automorphic representations of the unitary group*, arXiv: 1106.1127
- [42] T. Watson: *Rankin triple products and quantum chaos*, to appear in Ann. of Math. (2), Ph.D. dissertation, Princeton University, Princeton, 2002.

- [43] Xinyi Yuan, Shou-Wu Zhang, Wei Zhang. *The Gross–Zagier formula on Shimura curves*. Ann. of Math. Studies #184, ISBN: 9780691155920.
- [44] Zhiwei Yun: *The fundamental lemma of Jacquet and Rallis*. With an appendix by Julia Gordon. Duke Math. J. 156 (2011), no. 2, 167227.
- [45] Zhang, Shouwu. *Heights of Heegner points on Shimura curves*. Ann. of Math. (2) 153 (2001), no. 1, 27147
- [46] Zhang, Shou-Wu. *Gross-Zagier formula for $GL(2)$* . Asian J. Math. 5 (2001), no. 2, 183290.
- [47] Shouwu Zhang. *Gross-Zagier formula for $GL(2)$. II. Heegner points and Rankin L-series*, 191214, Math. Sci. Res. Inst. Publ., 49, Cambridge Univ. Press, Cambridge, 2004.
- [48] Shouwu Zhang. *Linear forms, algebraic cycles, and derivatives of L-series*, preprint.
- [49] W. Zhang. *On arithmetic fundamental lemmas*, Invent. Math., Volume 188, Number 1 (2012), 197-252.
- [50] W. Zhang. *Fourier transform and the global Gan–Gross–Prasad conjecture for unitary groups*, Preprint, 2011. Available at <http://www.math.columbia.edu/~wzhang/math/online/transfer.pdf>

Wei Zhang: Department of Mathematics, Columbia University, MC 4423 2990 Broadway, New York, NY 10027.

Email address: `wzhang@math.columbia.edu`.